

Prof. Arkady Leiderman

Fundamentals of Analysis for EE

Homework 2

Question 1. Consider the following subset of the plane:

$$A = \{(x, \sin(\frac{1}{x})) : x > 0\} \subset \mathbf{R}^2. \text{ What is the closure } \text{cl}(A) \text{ in } \mathbf{R}^2?$$

Question 2. Let $f(x) : \mathbf{R} \rightarrow \mathbf{R}$ be a function.

(a) Prove that if $f(x)$ is continuous at every point x , then the graph of the function

$$\Gamma = \{(x, f(x)) : x \in \mathbf{R}\} \text{ is a closed subset of } \mathbf{R}^2.$$

(b) Is the converse also true: If the graph of a function is closed in \mathbf{R}^2 , then the function $f(x)$ is continuous at every point?

Question 3. Let (X, d) be a metric space and A be a non-empty subset of X .

Define a number $\rho(x, A) = \inf\{d(x, y) : y \in A\}$ for every $x \in X$.

Prove that the closure $\text{cl}(A) = \{x \in X : \rho(x, A) = 0\}$.

Question 4. A metric space (X, d) is called separable if there is a countable set $A \subseteq X$ such that $\text{cl}(A) = X$. A family of open sets $B = \{U_\alpha\}$, where every $U_\alpha \subseteq X$, is called a base for X if for every point $x \in X$ and every open set V containing x there is $U_\alpha \in B$ such that $x \in U_\alpha \subseteq V$.

(a) Prove that any Euclidean space \mathbf{R}^n is separable.

(b) Prove that any metric space (X, d) is separable if and only if it has a countable base.

Question 5. In $X = \mathbf{R}$ define $d(x, y) = |\text{arctg}(x) - \text{arctg}(y)|$ for every $x, y \in X$.

(a) Prove that $d(x, y)$ defines a metric in X .

(b) Is (X, d) a complete metric space?

Question 6. Let $C[-1,1]$ be a metric space of continuous functions with the metric

$$d(f, g) = \int_{-1}^1 |f(x) - g(x)| dx. \text{ Consider the following sequence } \{f_n(x)\}_{n=1}^{\infty} :$$

$$f_n(x) = \begin{cases} 1 & \text{if } x \in [1/n, 1] \\ nx & \text{if } x \in [-1/n, 1/n] \\ -1 & \text{if } x \in [-1, -1/n] \end{cases}$$

Prove that $\{f_n(x)\}_{n=1}^{\infty}$ is a Cauchy sequence. Is the metric d complete?

Question 7. Let (X, d_X) and (Y, d_Y) be two metric spaces.

(a) Assume that a mapping $f : (X, d_X) \rightarrow (Y, d_Y)$ satisfies the Lipschitz condition:

$$\exists K > 0 \forall x_1, x_2 \in X \quad d_Y(f(x_1), f(x_2)) \leq K d_X(x_1, x_2).$$

Prove that $f(x)$ is a uniformly continuous mapping on (X, d_X) .

(b) Fix a point $p \in X$ and define a function $f : X \rightarrow \mathbf{R}$ by the formula: $f(x) = d_X(x, p)$.

Prove that $f(x)$ is a uniformly continuous function on (X, d_X) .

Question 8. Consider a closed ball in the space $C[0,1]$:

$$\bar{B} = \{f \in C[0,1] : \max_{x \in [0,1]} |f(x)| \leq 1\}. \text{ Is the set } \bar{B} \text{ compact?}$$

Question 9. Prove that if (X, d) is a compact metric space, then d is a complete metric.

Question 10. Let (X, d_X) and (Y, d_Y) be two metric spaces and $f : (X, d_X) \rightarrow (Y, d_Y)$

be a continuous mapping.

(a) Prove that if $K \subset X$ is a compact set, then its image $f(K) \subset Y$ is also a compact set.

(b) Let (X, d_X) be the Euclidean space \mathbf{R}^n . Prove that if $A \subset \mathbf{R}^n$ is a bounded set, then its image $f(A) \subset Y$ is also a bounded set.