

המחלקה למתמטיקה, בן-גוריון

---

---

## קולוקוויום

---

---

ביום שלישי, 26 בנובמבר, 2019

בשעה 14:30 – 15:30

ב-101 Math

ההרצאה

**On bounded continuous solutions of the  
rescaled archetypal equation**

תינתן על-ידי

Gregory Defel (BGU)

תקציר:

# On bounded continuous solutions of the archetypal equation with rescaling

Gregory Derfel

We'll start from a brief general introduction in equations with rescaling, that does not require any prerequisites.

Then we turn to the problem indicated in the title. Namely, we study the “archetypal functional equation  $y(x) = \int_{R^2} y(a(x-b)) \mu(da, db)$  ( $x \in R$ ), equivalently,  $y(x) = E\{y(\alpha(x-\beta))\}$ , where  $E$  is expectation with respect to the distribution  $\mu$  of random coefficients  $(\alpha, \beta)$ .

Particular cases include: (i)  $y(x) = \sum_i p_i y(a_i(x-b_i))$  and (ii)  $y'(x) + y(x) = \sum_i p_i y(a_i(x-b_i))$  (pantograph equation), both subject to the balance condition  $\sum_i p_i = 1$  ( $p_i > 0$ ).

Existence of non-trivial (i.e. non-constant) bounded continuous solutions is governed by the value  $K := \int_{R^2} \ln |a| \mu(da, db) = E\{\ln |\alpha|\}$ ; namely, under mild technical conditions no such solutions exist whenever  $K < 0$ , whereas if  $K > 0$  (and  $\alpha > 0$ ) then there is a non-trivial solution

In the critical case  $K = 0$ , we prove a Liouville theorem subject to the uniform continuity of  $y(\cdot)$ . The latter is guaranteed under a mild regularity assumption on the density of  $\beta$  conditioned on  $\alpha$ , which is satisfied for a large class of examples including the pantograph equation (ii).

Further results are obtained in the supercritical case  $K > 0$ , including existence, uniqueness and a maximum principle. The case with  $P(\alpha < 0) > 0$  is drastically different from that with  $\alpha > 0$ ; in particular, we prove that a bounded solution  $y(\cdot)$  possessing limits at  $\pm\infty$  must be constant.

The proofs employ martingale techniques applied to the martingale  $y(X_n)$ , where  $(X_n)$  is an associated Markov chain with jumps of the form  $x \rightarrow \alpha(x - \beta)$ .