המחלקה למתמטיקה, בן-גוריון

גאומטריה אלגברית ותורת המספרים

ביום רביעי, 26 באפריל, 2017

16:30 – 15:10 בשעה

101- Math**i**

ההרצאה

unramfiied the and Obstructions Homotopical problem Galois Inverse

תינתן על-ידי

(HU) Schlank Tomer

תקציר:



Ben Gurion University - Mathematics Algebraic Geometry and Number Theory Seminar

Speaker	Tomer Schlank (HU)
Title	Homotopical Obstructions and the unramified Inverse Galois problem
Date	Wednesday, 26 April 2017
Time	15:10 – 16:30 (starts 15:10 sharp)
Location	Room -101 in Building 58

Given a number field K, the unramified Inverse Galois problem is concerned with the question which finite groups \$G\$ can be realized as Galois groups of Galois unramified extensions \$L/K\$. The two main ways to attack the problem is by using class field theory (to analyze solvable extensions) and discriminant bounds (to analyze Fields \$K\$ of small discriminant). The goal of this talk is to show how using homotopical methods one can get results in the non-solvable case with no bound on the discriminant. We will begin by describing a general method to obtain homotopy theoretical obstructions to problems in Galois theory called "Embedding problems". Then we will explain how Abstract to employ these obstructions to study the unramified inverse Galois problem. Specifically, using these obstructions on embedding problems with a non-solvable kernel, we'll give an example of an infinite family of groups {G i}i together with an infinite family of quadratic number fields such that for any number field K in this family, the maximal solvable quotient of G i is realizable as an unramified Galois group over K; but G i itself is not.

This is a joint work with Magnus Carlson

(updated 6 Apr 2017)