



# ***Variation of complex and $p$ -adic regulators***

Amnon Besser

- Work of Rodriguez Villegas on “modular mahler measures” suggests that the derivative of the regulator in families is much simpler than the regulator itself, depending only on the de Rham Chern class.

- Work of Rodriguez Villegas on “modular mahler measures” suggests that the derivative of the regulator in families is much simpler than the regulator itself, depending only on the de Rham Chern class.
- This fact, in different guises, has been observed by various people.

- Work of Rodriguez Villegas on “modular mahler measures” suggests that the derivative of the regulator in families is much simpler than the regulator itself, depending only on the de Rham Chern class.
- This fact, in different guises, has been observed by various people.
- The proof depends on a simple homological algebra Lemma, due to Jannsen.

- Work of Rodriguez Villegas on “modular mahler measures” suggests that the derivative of the regulator in families is much simpler than the regulator itself, depending only on the de Rham Chern class.
- This fact, in different guises, has been observed by various people.
- The proof depends on a simple homological algebra Lemma, due to Jannsen.
- It could explain various instances of polylogarithms

- Work of Rodriguez Villegas on “modular mahler measures” suggests that the derivative of the regulator in families is much simpler than the regulator itself, depending only on the de Rham Chern class.
- This fact, in different guises, has been observed by various people.
- The proof depends on a simple homological algebra Lemma, due to Jannsen.
- It could explain various instances of polylogarithms
- It also has (conjecturally) a  $p$ -adic analogue.

# ***Mahler measures as regulators***

---

$$m(P) = \frac{1}{(2\pi i)^n} \int_{T^n} \log |P(z_1, \dots, z_n)| \frac{dz_1}{z_1} \cdots \frac{dz_n}{z_n}$$

# ***Mahler measures as regulators***

---

$$m(P) = \frac{1}{(2\pi i)^n} \int_{T^n} \log |P(z_1, \dots, z_n)| \frac{dz_1}{z_1} \dots \frac{dz_n}{z_n}$$

Deninger (Journal of the AMS 10 259-281 (1997)):

Take  $Z = \{P(z_1, \dots, z_n) = 0\}$ ,  $X = \{z \neq 0\}^n - Z$



# *Mahler measures as regulators*

$$m(P) = \frac{1}{(2\pi i)^n} \int_{T^n} \log |P(z_1, \dots, z_n)| \frac{dz_1}{z_1} \dots \frac{dz_n}{z_n}$$

Deninger (Journal of the AMS 10 259-281 (1997)):

Take  $Z = \{P(z_1, \dots, z_n) = 0\}$ ,  $X = \{z \neq 0\}^n - Z$

$$\begin{aligned} K_{n+1}(X) &\xrightarrow{r_D} H_D^{n+1}(X, \mathbb{R}(n+1)) \\ &= H^n(X, (2\pi i)^n \mathbb{R}) \xrightarrow{(2\pi i)^{-n} \int_{T^n}} \mathbb{R} \end{aligned}$$

# Mahler measures as regulators

$$m(P) = \frac{1}{(2\pi i)^n} \int_{T^n} \log |P(z_1, \dots, z_n)| \frac{dz_1}{z_1} \cdots \frac{dz_n}{z_n}$$

Deninger (Journal of the AMS 10 259-281 (1997)):

Take  $Z = \{P(z_1, \dots, z_n) = 0\}$ ,  $X = \{z \neq 0\}^n - Z$

$$\begin{aligned} K_{n+1}(X) &\xrightarrow{r_D} H_D^{n+1}(X, \mathbb{R}(n+1)) \\ &= H^n(X, (2\pi i)^n \mathbb{R}) \xrightarrow{(2\pi i)^{-n} \int_{T^n}} \mathbb{R} \end{aligned}$$

Deninger computes:

$$\{P, z_1, \dots, z_n\} := P \wedge z_1 \wedge \cdots \wedge z_n \mapsto m(P)$$

He further shows that Mahler measures, or differences of Mahler measures, can sometimes be related to elements in the K-theory of  $Z$ , hence via the Beilinson conjectures to special values of  $L$ -functions.

# ***Modular Mahler measures***

---

Rodriguez Villegas (Topics in number theory 17-48)  
Mahler measures in families:

$$P_k(x, y) = k - P(x, y)$$

# ***Modular Mahler measures***

---

Rodriguez Villegas (Topics in number theory 17-48)  
Mahler measures in families:

$$P_k(x, y) = k - P(x, y)$$

$$m(P_k) = -\operatorname{Re}(\tilde{m}(1/k)), \text{ where}$$

$$\tilde{m}(s) = \log(s) - \frac{1}{(2\pi i)^2} \int_{T^2} \log(1 - sP(x, y)) \frac{dx}{x} \frac{dy}{y}$$

# Modular Mahler measures

Rodriguez Villegas (Topics in number theory 17-48)  
Mahler measures in families:

$$P_k(x, y) = k - P(x, y)$$

$$m(P_k) = -\operatorname{Re}(\tilde{m}(1/k)), \text{ where}$$

$$\tilde{m}(s) = \log(s) - \frac{1}{(2\pi i)^2} \int_{T^2} \log(1 - sP(x, y)) \frac{dx}{x} \frac{dy}{y}$$

and then he find that  $d\tilde{m}(s) = u_0(s)ds/s$  where  $u_0$  has (sometimes) really nice properties:

# Modular Mahler measures

Rodriguez Villegas (Topics in number theory 17-48)  
Mahler measures in families:

$$P_k(x, y) = k - P(x, y)$$

$$m(P_k) = -\operatorname{Re}(\tilde{m}(1/k)), \text{ where}$$

$$\tilde{m}(s) = \log(s) - \frac{1}{(2\pi i)^2} \int_{T^2} \log(1 - sP(x, y)) \frac{dx}{x} \frac{dy}{y}$$

and then he find that  $d\tilde{m}(s) = u_0(s)ds/s$  where  $u_0$  has (sometimes) really nice properties:

- It satisfies a Picard-Fuchs equation

# Modular Mahler measures

Rodriguez Villegas (Topics in number theory 17-48)  
Mahler measures in families:

$$P_k(x, y) = k - P(x, y)$$

$$m(P_k) = -\operatorname{Re}(\tilde{m}(1/k)), \text{ where}$$

$$\tilde{m}(s) = \log(s) - \frac{1}{(2\pi i)^2} \int_{T^2} \log(1 - sP(x, y)) \frac{dx}{x} \frac{dy}{y}$$

and then he find that  $d\tilde{m}(s) = u_0(s)ds/s$  where  $u_0$  has (sometimes) really nice properties:

- It satisfies a Picard-Fuchs equation
- It is a modular form when properly interpreted



# Modular Mahler measures

Rodriguez Villegas (Topics in number theory 17-48)  
Mahler measures in families:

$$P_k(x, y) = k - P(x, y)$$

$$m(P_k) = -\operatorname{Re}(\tilde{m}(1/k)), \text{ where}$$

$$\tilde{m}(s) = \log(s) - \frac{1}{(2\pi i)^2} \int_{T^2} \log(1 - sP(x, y)) \frac{dx}{x} \frac{dy}{y}$$

and then he find that  $d\tilde{m}(s) = u_0(s)ds/s$  where  $u_0$  has (sometimes) really nice properties:

- It satisfies a Picard-Fuchs equation
- It is a modular form when properly interpreted

**Is there an explanation for this?**

# ***Integral Deligne cohomology***

---

Our first clue: Real Deligne cohomology is “the real part” of integral Deligne cohomology:

# *Integral Deligne cohomology*

---

Our first clue: Real Deligne cohomology is “the real part” of integral Deligne cohomology:  
In (almost) complete generality

$$H_D^{m+1}(X, \mathbb{Z}(k)) = \frac{H^m(X, \mathbb{C})}{F^k H^m(X, \mathbb{C}) + H^m(X, (2\pi i)^k \mathbb{Z})}$$

which is also known as an “intermediate jacobian”,  
where

$$F^k H^m(X, \mathbb{C}) = \bigoplus_{p \geq k} H^{p, m-p}$$

$H_D^{m+1}(X, \mathbb{R}(k))$  is the same thing with  $\mathbb{R}$  instead of  $\mathbb{Z}$ .

For  $m = n$ ,  $k = n + 1$

$$H_D^{m+1}(X, \mathbb{Z}(k)) = \frac{H^m(X, \mathbb{C})}{F^k H^m(X, \mathbb{C}) + H^m(X, (2\pi i)^k \mathbb{Z})}$$

For  $m = n$ ,  $k = n + 1$

$$H_D^{n+1}(X, \mathbb{R}(n+1)) = \frac{H^n(X, \mathbb{C})}{F^{n+1}H^n(X, \mathbb{C}) + H^n(X, (2\pi i)^{n+1}\mathbb{R})}$$

For  $m = n$ ,  $k = n + 1$  there is no  $F^{n+1}H^n(X, \mathbb{C})$

$$H_D^{n+1}(X, \mathbb{R}(n+1)) = \frac{H^n(X, \mathbb{C})}{H^n(X, (2\pi i)^{n+1} \mathbb{R})}$$

For  $m = n$ ,  $k = n + 1$  there is no  $F^{n+1}H^n(X, \mathbb{C})$

$$H_D^{n+1}(X, \mathbb{R}(n+1)) = \frac{H^n(X, \mathbb{C})}{H^n(X, (2\pi i)^{n+1} \mathbb{R})}$$

and we get  $H^n(X, (2\pi i)^n \mathbb{R})$  by taking the real or imaginary value.

# ***Families of varieties and normal functions***

---

Consider a family of varieties  $X_s$   
 $X$  is the total space.



# *Families of varieties and normal functions*

---

Consider a family of varieties  $X_s$   
 $X$  is the total space.

Normal functions = nice functions  $f(s) \in H_D^{m+1}(X_s, \mathbb{Z}(k))$

# *Families of varieties and normal functions*

---

Consider a family of varieties  $X_s$   
 $X$  is the total space.

Normal functions = nice functions  $f(s) \in H_D^{m+1}(X_s, \mathbb{Z}(k))$

“Cycle people” know that a cycle  $Z$  on  $X$  gives a  
normal function  $f_Z(s) = r_D(Z \cap X_s)$

# *Families of varieties and normal functions*

---

Consider a family of varieties  $X_s$   
 $X$  is the total space.

Normal functions = nice functions  $f(s) \in H_D^{m+1}(X_s, \mathbb{Z}(k))$

“Cycle people” know that a cycle  $Z$  on  $X$  gives a normal function  $f_Z(s) = r_D(Z \cap X_s)$

... But others don't know that the same is true for K-theory:  $\alpha \in K_i(X)$  will give a normal function

$$f_\alpha(s) = r_D(\alpha|_{X_s})$$

# *Families of varieties and normal functions*

---

Consider a family of varieties  $X_s$   
 $X$  is the total space.

Normal functions = nice functions  $f(s) \in H_D^{m+1}(X_s, \mathbb{Z}(k))$

“Cycle people” know that a cycle  $Z$  on  $X$  gives a normal function  $f_Z(s) = r_D(Z \cap X_s)$

... But others don't know that the same is true for K-theory:  $\alpha \in K_i(X)$  will give a normal function

$$f_\alpha(s) = r_D(\alpha|_{X_s})$$

In particular, for a family of polynomials  $P_s$  we get a normal function  $f_P(s) = r_D(\{P_s, z_1, \dots, z_n\})$

## ***Enters Gauss-Manin***

---

On  $f_P(s)$  we want to  
first do  $\int_{T^n}$ , then differentiate

## ***Enters Gauss-Manin***

---

On  $f_P(s)$  we want to  
first do  $\int_{T^n}$ , then differentiate

It is well known that we can instead  
first apply the Gauss-Manin connection, then do  $\int_{T^n}$ .

On  $f_P(s)$  we want to  
first do  $\int_{T^n}$ , then differentiate

It is well known that we can instead  
first apply the Gauss-Manin connection, then do  $\int_{T^n}$ .

Gauss-Manin:  $\nabla : \mathcal{H}^m(X/S) \rightarrow \Omega^1(S, \mathcal{H}^m(X/S))$ .

On  $f_P(s)$  we want to  
first do  $\int_{T^n}$ , then differentiate

It is well known that we can instead  
first apply the Gauss-Manin connection, then do  $\int_{T^n}$ .

Gauss-Manin:  $\nabla : \mathcal{H}^m(X/S) \rightarrow \Omega^1(S, \mathcal{H}^m(X/S))$ .

To apply  $\nabla$  to  $f(s)$  first lift to  $\tilde{f}(s) \in \mathcal{H}^m(X/S)$ , then  
apply  $\nabla$ .



On  $f_P(s)$  we want to  
first do  $\int_{T^n}$ , then differentiate

It is well known that we can instead  
first apply the Gauss-Manin connection, then do  $\int_{T^n}$ .

Gauss-Manin:  $\nabla : \mathcal{H}^m(X/S) \rightarrow \Omega^1(S, \mathcal{H}^m(X/S))$ .

To apply  $\nabla$  to  $f(s)$  first lift to  $\tilde{f}(s) \in \mathcal{H}^m(X/S)$ , then  
apply  $\nabla$ .

The integral part disappears and we get

$$\nabla f(s) \in \Omega^1(S, \mathcal{H}^m(X/S)/F^{k-1}\mathcal{H}^m(X/S))$$

## ***The main theorem***

---

**Main Theorem:** for a class in K-theory  $\alpha$  the derivative of the normal function,  $\nabla f_\alpha(s)$ , can be expressed in terms of de Rham class  $ch(\alpha) \in F^k H^{m+1}(X)$ .

## ***The main theorem***

---

**Main Theorem:** for a class in K-theory  $\alpha$  the derivative of the normal function,  $\nabla f_\alpha(s)$ , can be expressed in terms of de Rham class  $ch(\alpha) \in F^k H^{m+1}(X)$ .  
More precisely, with respect to the Leray spectral sequence

$$H^i(S, \mathcal{H}^{m+1-i}(X/S)) \Rightarrow H^{m+1}(X)$$

$ch(\alpha)$  maps to 0 in  $H^0(S, \mathcal{H}^{m+1}(X/S))$  and  $-\nabla f_\alpha(s)$  gives its image in  $H^1(S, \mathcal{H}^m(X/S))$ .

## *The main theorem*

**Main Theorem:** for a class in K-theory  $\alpha$  the derivative of the normal function,  $\nabla f_\alpha(s)$ , can be expressed in terms of de Rham class  $ch(\alpha) \in F^k H^{m+1}(X)$ .  
More precisely, with respect to the Leray spectral sequence

$$H^i(S, \mathcal{H}^{m+1-i}(X/S)) \Rightarrow H^{m+1}(X)$$

$ch(\alpha)$  maps to 0 in  $H^0(S, \mathcal{H}^{m+1}(X/S))$  and  $-\nabla f_\alpha(s)$  gives its image in  $H^1(S, \mathcal{H}^m(X/S))$ .

Many people know about this in various forms: Voisin, Collino, Bloch, de Jeu, Saito ....

## *Why do we care?*

---

- Since the de Rham class is an algebraic object, it is much simpler than the Deligne cohomology class.

## *Why do we care?*

---

- Since the de Rham class is an algebraic object, it is much simpler than the Deligne cohomology class.
- It explains the results of Rodriguez Villegas.

## *Why do we care?*

---

- Since the de Rham class is an algebraic object, it is much simpler than the Deligne cohomology class.
- It explains the results of Rodriguez Villegas.
- Conjecturally, you can prove the same result with Deligne cohomology replaced by its  $p$ -adic analogue – Syntomic cohomology. Moreover, the original class can be reconstructed by using a Coleman integral.

## *Why do we care?*

---

- Since the de Rham class is an algebraic object, it is much simpler than the Deligne cohomology class.
- It explains the results of Rodriguez Villegas.
- Conjecturally, you can prove the same result with Deligne cohomology replaced by its  $p$ -adic analogue – Syntomic cohomology. Moreover, the original class can be reconstructed by using a Coleman integral.
- It can be used to compute regulators in many interesting cases, in particular related to many types of polylogarithms.



## Homological algebra Lemma (Jannsen, B.)

$$\begin{array}{ccccc} X & \xrightarrow{\alpha} & Y & \xrightarrow{\beta} & C(X \rightarrow Y) \\ \varepsilon \downarrow & & \phi \downarrow & & \downarrow \\ Z & \xrightarrow{\gamma} & W & \longrightarrow & C(Z \rightarrow W) \\ \eta \downarrow & & \downarrow & & \downarrow \\ C(X \rightarrow Z) & \longrightarrow & C(Y \rightarrow W) & \longrightarrow & U \end{array}$$

leading to

$$\begin{array}{ccccccc}
 H_{X,Z}^{i-1} & \longrightarrow & H_{Y,W}^{i-1} & \longrightarrow & H_U^{i-1} & \longrightarrow & H_{X,Z}^i \\
 \downarrow & & \downarrow & & \downarrow & & \downarrow \\
 H_X^i & \longrightarrow & H_Y^i & \longrightarrow & H_{X,Y}^i & \longrightarrow & H_X^{i+1} \\
 \downarrow & & \downarrow & & \downarrow & & \downarrow \\
 H_Z^i & \longrightarrow & H_W^i & \longrightarrow & H_{Z,W}^i & \longrightarrow & H_Z^{i+1} \\
 \downarrow & & \downarrow & & \downarrow & & \downarrow \\
 H_{X,Z}^i & \longrightarrow & H_{Y,W}^i & \longrightarrow & H_U^i & \longrightarrow & H_{X,Z}^{i+1}
 \end{array}$$

Theorem follows from applying this to the following diagram

$$\begin{array}{ccccc}
 ?? & \longrightarrow & \mathbb{Z}_{\mathcal{D}}(k)_X & \longrightarrow & \mathbb{Z}_{\mathcal{D}}(k)_{X/S} \\
 \downarrow & & \downarrow & & \downarrow \\
 0 \oplus \Omega_S^1 \otimes F^k \Omega_X^\bullet[-1] & \longrightarrow & \mathbb{Z} \oplus F^k \Omega_X^\bullet & \longrightarrow & \mathbb{Z} \oplus F^k \Omega_{X/S}^\bullet \\
 \downarrow & & \downarrow & & \downarrow \\
 \Omega_S^1 \otimes \Omega_X^\bullet[-1] & \longrightarrow & \Omega_X^\bullet & \longrightarrow & \Omega_{X/S}^\bullet
 \end{array}$$

## ***$p$ -adic Mahler measures I***

---

We want a  $p$ -adic analogue of Mahler measures.  
From the regulator point of view we need two things:

# ***$p$ -adic Mahler measures I***

---

We want a  $p$ -adic analogue of Mahler measures.  
From the regulator point of view we need two things:

- An analogue of the regulator – The syntomic regulator

# ***p-adic Mahler measures I***

We want a  $p$ -adic analogue of Mahler measures.  
From the regulator point of view we need two things:

- An analogue of the regulator – The syntomic regulator
- An analogue of integration against  $T^n$  – Schnirelman integral

$$\int_{T_p^n} f(z_1, \dots, z_n) \frac{dz_1}{z_1} \cdots \frac{dz_n}{z_n} := \lim_{\substack{N \rightarrow \infty \\ (N, p) = 1}} \frac{1}{N^n} \sum_{\zeta \in \mu_N^n} f(\zeta)$$

We obtain the  $p$ -adic Mahler measure

$$m_p(P) = \int_{T_p^n} \log(P(z_1, \dots, z_n)) \frac{dz_1}{z_1} \cdots \frac{dz_n}{z_n}$$

We obtain the  $p$ -adic Mahler measure

$$m_p(P) = \int_{T_p^n} \log(P(z_1, \dots, z_n)) \frac{dz_1}{z_1} \dots \frac{dz_n}{z_n}$$

In the case  $n = 1$  this gives for  $P = a_m t^m + \dots + a_r t^r$

$$m_p(P) = \log_p(a_m) + \sum_{|\alpha|_p > 1} \log_p(\alpha)$$



## ***$p$ -adic Mahler measures II***

---

Instead of using Schnirelman integral we import  $T^n$  into the  $p$ -adic world.

## ***$p$ -adic Mahler measures II***

---

Instead of using Schnirelman integral we import  $T^n$  into the  $p$ -adic world. This requires

- Extending scalars to  $B_{dR}$

## ***$p$ -adic Mahler measures II***

---

Instead of using Schnirelman integral we import  $T^n$  into the  $p$ -adic world. This requires

- Extending scalars to  $B_{dR}$
- Fixing embeddings of  $\bar{\mathbb{Q}}$  to  $\mathbb{C}$  and to  $\mathbb{C}_p$ , collectively called  $\sigma$ .

## ***$p$ -adic Mahler measures II***

---

Instead of using Schnirelman integral we import  $T^n$  into the  $p$ -adic world. This requires

- Extending scalars to  $B_{dR}$
- Fixing embeddings of  $\bar{\mathbb{Q}}$  to  $\mathbb{C}$  and to  $\mathbb{C}_p$ , collectively called  $\sigma$ .

This gives  $m_{p,\sigma}(P)$ .

## ***$p$ -adic Mahler measures II***

Instead of using Schnirelman integral we import  $T^n$  into the  $p$ -adic world. This requires

- Extending scalars to  $B_{dR}$
- Fixing embeddings of  $\bar{\mathbb{Q}}$  to  $\mathbb{C}$  and to  $\mathbb{C}_p$ , collectively called  $\sigma$ .

This gives  $m_{p,\sigma}(P)$ .

In the case  $n = 1$  this gives for  $P = a_m t^m + \cdots a_r t^r$

$$m_{p,\sigma}(P) = \log_p(a_m) + \sum_{|\alpha|_\infty > 1} \log_p(\alpha)$$

## ***$p$ -adic Mahler measures II***

Instead of using Schnirelman integral we import  $T^n$  into the  $p$ -adic world. This requires

- Extending scalars to  $B_{dR}$
- Fixing embeddings of  $\bar{\mathbb{Q}}$  to  $\mathbb{C}$  and to  $\mathbb{C}_p$ , collectively called  $\sigma$ .

This gives  $m_{p,\sigma}(P)$ .

In the case  $n = 1$  this gives for  $P = a_m t^m + \cdots a_r t^r$

$$m_{p,\sigma}(P) = \log_p(a_m) + \sum_{|\alpha|_\infty > 1} \log_p(\alpha)$$

This can sometimes be tied to special values of  $p$ -adic  $L$ -functions.