

# Ordinary Differential Equations for Math

(201.1.0061. Spring 2023. Dmitry Kerner)

## Homework 1. Submission date: 23.03.2023

Questions to submit (by mail): 1 b. 1 d. 2 c. 3 a. 3 b. 4 b.

Homeworks must be either typed (in Latex) or written in readable handwriting and scanned in readable resolution.



1.
  - a. The earth is defined by the condition  $\{\underline{x} \mid x_n \leq 0\} \subset \mathbb{R}^n$ . Suppose every freely falling particle (with  $x_n > 0$ ) experiences the constant acceleration,  $\vec{a} = -g \cdot \hat{x}_n$ . (The law of Galileo,  $g = 9.8m/sec^2$ .) A frog jumps at the moment  $t_0$  from the point  $0 \in \mathbb{R}^n$ , with the initial velocity  $\vec{v}$  (here  $v_n > 0$ ). Find the trajectory  $\underline{x}(t)$ , the total time lapse of the jump (i.e.  $t_1 - t_0$ ) and the total displacement (i.e.  $\underline{x}(t_1) - \underline{x}(t_0)$ ).
  - b. A particle moves on the standard sphere  $\mathbb{S}^{n-1} \subset \mathbb{R}^n$ . Prove:  $\vec{v}(t) \perp \underline{x}(t)$  for each  $t$ . Suppose  $n = 2$  and  $\|\vec{v}\| = const$ . Prove:  $\vec{a}(t) \parallel \underline{x}(t)$  for each  $t$ . (wiki: centrifugal force)
  - c. How does a differentiable coordinate change,  $\mathbb{R}^n \xrightarrow{\phi} \mathbb{R}^n$ ,  $\underline{x} \rightarrow \phi(\underline{x})$ , affect  $\vec{v}$ ,  $\vec{a}$ ?
  - d. Take the second law of Newton in dimension 1, i.e.  $m \cdot \vec{a} = \vec{F}(x)$ , for a continuous vector field  $\mathbb{R}^1 \xrightarrow{\vec{F}} \mathbb{R}^1$ . Multiply this 2'nd order ODE by  $\vec{v}$  and integrate. You get a 1'st order ODE, called the "Energy conservation law". Here  $\frac{m \cdot \vec{v}^2}{2}$  is the kinetic energy, while  $-\int F(x)dx$  is the potential energy.  
(To extend this to  $n > 1$  one needs some ingredients from Geometric Calculus 2.)
  
2. Consider the equation  $x' + f(t) \cdot x = g(t)$ , for  $f, g \in C^0(a, b)$ .
  - a. We have obtained the complete solution in the class. Rederive the general formula. What is the dimensionality of the family of solutions? (It seems to depend on two constants?)
  - b. Impose the initial condition  $x(t_0) = x_0$ . Prove: the solution is unique and is globally defined.
  - c. For which pairs  $(t_0, x_0) \in \mathbb{R}^2$  does the IVP  $t \cdot x' + x = \sin(t)$ ,  $x(t_0) = x_0$  posses a solution? When is it unique? Draw the field of directions and the integral curves.
  
3. The hare runs over  $\mathbb{R}_{x,y}^2$  with constant velocity  $(0, v_h)$ , starting at  $t = 0$  from the origin. The dog starts to run from the point  $(x_0, 0)$  at  $t = 0$  with velocity  $\vec{v}$  satisfying:  $\|\vec{v}\| = v_d$ , and at each moment  $\vec{v}$  is directed towards the hare. Present the trajectory of the dog as  $y = y(x)$ .
  - a. Prove: the function  $y(x)$  satisfies the equation  $x \cdot y'' = \frac{v_h}{v_d} \cdot \sqrt{1 + (y')^2}$ .  
(Hint: write all the equations on the functions  $x(t), y(t)$ . Observe that these functions are invertible. Exclude  $t$  from this system of equations.)
  - b. Substitute  $z = y'$  to convert this 2'nd order equation to a first order ODE. Solve it.
  - c. Give the necessary and sufficient condition (on  $v_h, v_d$ ) for the dog to catch the hare in finite time. (Verify that the natural guess means the converges of some improper integral.)
  - d. Wiki: *The curve of pursuit*. (The french version is better than the english one, 13.03.2023.)
  
4. Suppose  $x(t)$  is a solution of the equation  $x' = f(t, x)$ , for some  $f \in C^\infty(\mathcal{U})$ .
  - a. Prove: if  $t_0$  is an extremum of the function  $x(t)$  then  $f(t_0, x(t_0)) = 0$ .  
Is this also a sufficient condition for an extremum of the solution?
  - b. Suppose  $f(t_0, x(t_0)) = 0$  and  $\partial_t f|_{(t_0, x(t_0))} > 0$ . Prove:  $x(t)$  has a local minimum at  $t_0$ .
  - c. Suppose  $\partial_t^j f|_{(t_0, x(t_0))} = 0$  for  $j = 0, \dots, r-1$  and  $\partial_t^r f|_{(t_0, x(t_0))} < 0$ . Is  $t_0$  a local min/max of  $x(t)$ ?