

ON THE \mathbf{cd} -INDEX AND γ -VECTOR OF S^* -SHELLABLE CW-SPHERES

SATOSHI MURAI AND ERAN NEVO

ABSTRACT. We show that the γ -vector of the order complex of any polytope is the f -vector of a balanced simplicial complex. This is done by proving this statement for a subclass of Stanley’s S -shellable spheres which includes all polytopes. The proof shows that certain parts of the \mathbf{cd} -index, when specializing $\mathbf{c} = 1$ and considering the resulted polynomial in \mathbf{d} , are the f -polynomials of simplicial complexes that can be colored with “few” colors. We conjecture that the \mathbf{cd} -index of a regular CW-sphere is itself the *flag* f -vector of a colored simplicial complex in a certain sense.

1. INTRODUCTION

Let P be an $(n-1)$ -dimensional regular CW-sphere (that is, a regular CW-complex which is homeomorphic to an $(n-1)$ -dimensional sphere). In face enumeration, one of the most important combinatorial invariants of P is the \mathbf{cd} -index. The \mathbf{cd} -index $\Phi_P(\mathbf{c}, \mathbf{d})$ of P is a non-commutative polynomial in the variables \mathbf{c} and \mathbf{d} that encodes the flag f -vector of P . By the result of Stanley [St1] and Karu [Ka], it is known that the \mathbf{cd} -index $\Phi_P(\mathbf{c}, \mathbf{d})$ has non-negative integer coefficients. On the other hand, a characterization of the possible \mathbf{cd} -indices for regular CW-spheres, or other related families, e.g Gorenstien* posets, is still beyond reach. In this paper we take a step in this direction and establish some non-trivial upper bounds, as we detail now.

If we substitute 1 for \mathbf{c} in $\Phi_P(\mathbf{c}, \mathbf{d})$, we obtain a polynomial of the form

$$\Phi_P(1, \mathbf{d}) = \delta_0 + \delta_1 \mathbf{d} + \cdots + \delta_{\lfloor \frac{n}{2} \rfloor} \mathbf{d}^{\lfloor \frac{n}{2} \rfloor},$$

where $\lfloor \frac{n}{2} \rfloor$ is the integer part of $\frac{n}{2}$, such that each δ_i is a non-negative integer. In other words, δ_i is the sum of coefficients of monomials in $\Phi_P(\mathbf{c}, \mathbf{d})$ for which \mathbf{d} appears i times.

Let Δ be a (finite abstract) simplicial complex on the vertex set V . We say that Δ is k -colored if there is a map $c : V \rightarrow [k] = \{1, 2, \dots, k\}$, called a k -coloring map of Δ , such that if $\{x, y\}$ is an edge of Δ then $c(x) \neq c(y)$. Let $f_i(\Delta)$ denote the number of elements $F \in \Delta$ having cardinality $i + 1$, where $f_{-1}(\Delta) = 1$. The main result of this paper is the following.

Theorem 1.1. *Let P be an $(n-1)$ -dimensional S^* -shellable regular CW-sphere, and let $\Phi_P(1, \mathbf{d}) = \delta_0 + \delta_1 \mathbf{d} + \cdots + \delta_{\lfloor \frac{n}{2} \rfloor} \mathbf{d}^{\lfloor \frac{n}{2} \rfloor}$. Then there exists an $\lfloor \frac{n}{2} \rfloor$ -colored simplicial complex Δ such that*

$$\delta_i = f_{i-1}(\Delta) \quad \text{for } i = 0, 1, \dots, \lfloor \frac{n}{2} \rfloor.$$

Key words and phrases. polytope, γ -vector, \mathbf{cd} -index, balanced simplicial complex.

Research of the first author was partially supported by KAKENHI 22740018. Research of the second author was partially supported by an NSF Award DMS-0757828.

The precise definition of the S^* -shellability is given in Section 2. The most important class of S^* -shellable CW-spheres are the boundary complexes of polytopes. By the Kruskal-Katona Theorem (see e.g. [St2, II, Theorem 2.1]), the above theorem gives certain upper bound on δ_i in terms of δ_{i-1} . Better upper bounds are given by Frankl-Füredi-Kalai theorem which characterizes the f -vectors of k -colored complexes [FFK].

The numbers $\delta_0, \delta_1, \delta_2, \dots$ relate to the γ -vector (see Section 4 for the definition) of the barycentric subdivision (order complex) of P , namely the simplicial complex whose elements are the chains of nonempty cells in P ordered by inclusion. Indeed, as an application of Theorem 1.1 we prove the following.

Theorem 1.2. *Let P be an $(n - 1)$ -dimensional S^* -shellable regular CW-sphere and let $\text{sd}(P)$ be the barycentric subdivision of P . Then there exists an $\lfloor \frac{n}{2} \rfloor$ -colored simplicial complex Γ such that*

$$\gamma_i(\text{sd}(P)) = f_{i-1}(\Gamma) \quad \text{for } i = 0, 1, \dots, \lfloor \frac{n}{2} \rfloor.$$

Recall that an $(n - 1)$ -dimensional simplicial complex is said to be *balanced* if it is n -colored. If P is the boundary complex of an arbitrary convex n -dimensional polytope, then $\delta_{\lfloor \frac{n}{2} \rfloor}(P) > 0$ and we conclude the following.

Corollary 1.3. *Let P be the boundary complex of an n -dimensional polytope. Then the γ -vector of $\text{sd}(P)$ is the f -vector of a balanced simplicial complex.*

The above corollary supports the conjecture of Nevo and Petersen [NP, Conjecture 6.3] which states that the γ -vector of a flag homology sphere is the f -vector of a balanced simplicial complex. This conjecture was verified for the barycentric subdivision of simplicial homology spheres (in this case all the cells are simplices) in [NPT].

It would be natural to ask if the above theorems hold for all regular CW-spheres (or more generally, Gorenstein* posets). We conjecture a stronger statement on the \mathbf{cd} -index, see Conjecture 4.3.

This paper is organized as follows: in Section 2 we recall some known results on the \mathbf{cd} -index and define S^* -shellability, in Section 3 we prove our main theorem, Theorem 1.1, in Section 4 we derive consequences for γ -vectors and present a conjecture on the \mathbf{cd} -index, Conjecture 4.3.

2. \mathbf{cd} -INDEX OF S^* -SHELLABLE CW-SPHERES

In this section we recall some known results on the \mathbf{cd} -index.

Let P be a graded poset of rank $n + 1$ with the minimal element $\hat{0}$ and the maximal element $\hat{1}$. Let ρ denote the rank function of P . For $S \subset [n] = \{1, 2, \dots, n\}$, a chain $\hat{0} = \sigma_0 < \sigma_1 < \sigma_2 < \dots < \sigma_{k+1} = \hat{1}$ of P is called an S -flag if $\{\rho(\sigma_1), \dots, \rho(\sigma_k)\} = S$. Let $f_S(P)$ be the number of S -flags of P . Define $h_S(P)$ by

$$h_S(P) = \sum_{T \subset S} (-1)^{|S| - |T|} f_T(P),$$

where $|X|$ denotes the cardinality of a finite set X . The vectors $(f_S(P) : S \subset [n])$ and $(h_S(P) : S \subset [n])$ are called the *flag f -vector* and *flag h -vector* of P respectively.

Now we recall the definition of the \mathbf{cd} -index. For $S \subset [n]$, we define a non-commutative monomial $u_S = u_1 u_2 \cdots u_n$ in variables \mathbf{a} and \mathbf{b} by $u_i = \mathbf{a}$ if $i \notin S$ and $u_i = \mathbf{b}$ if $i \in S$. Let

$$\Psi_P(\mathbf{a}, \mathbf{b}) = \sum_{S \subset [n]} h_P(S) u_S.$$

For a graded poset P , let $\text{sd}(P)$ be the order complex of $P - \{\hat{0}, \hat{1}\}$. Thus

$$\text{sd}(P) = \{\{\sigma_1, \sigma_2, \dots, \sigma_k\} \subset P - \{\hat{0}, \hat{1}\} : \sigma_1 < \sigma_2 < \dots < \sigma_k\}.$$

We say that P is *Gorenstein** if the simplicial complex $\text{sd}(P)$ is a homology sphere. It is known that if P is Gorenstein* then $\Psi_P(\mathbf{a}, \mathbf{b})$ can be written as a polynomial $\Phi_P(\mathbf{c}, \mathbf{d})$ in $\mathbf{c} = \mathbf{a} + \mathbf{b}$ and $\mathbf{d} = \mathbf{ab} + \mathbf{ba}$ [BK], and this non-commutative polynomial $\Phi_P(\mathbf{c}, \mathbf{d})$ is called the \mathbf{cd} -index of P . Moreover, by the celebrated results due to Stanley [St1] (for convex polytopes) and Karu [Ka] (for Gorenstein* posets), the coefficients of $\Phi_P(\mathbf{c}, \mathbf{d})$ are non-negative integers.

Next, we define S^* -shellability of regular CW-spheres by slightly modifying the definition of S -shellability introduced by Stanley [St1, Definition 2.1].

Let P be a regular CW-sphere (a regular CW-complex which is homeomorphic to a sphere) and $\mathcal{F}(P)$ its face poset. Then the order complex of $\mathcal{F}(P)$ is a triangulation of a sphere, so the poset $\mathcal{F}(P) \cup \{\hat{0}, \hat{1}\}$ is Gorenstein*. We define the \mathbf{cd} -index of P by $\Phi_P(\mathbf{c}, \mathbf{d}) = \Phi_{\mathcal{F}(P) \cup \{\hat{0}, \hat{1}\}}(\mathbf{c}, \mathbf{d})$. For any cell σ of P , we write $\bar{\sigma}$ for the closure of σ . For an $(n-1)$ -dimensional regular CW-sphere P , let ΣP be the suspension of P , in other words, ΣP is the n -dimensional regular CW-sphere obtained from P by attaching two n -dimensional cells τ_1 and τ_2 such that $\partial\tau_1 = \partial\tau_2 = P$. Also, for an $(n-1)$ -dimensional regular CW-ball P (a regular CW-complex which is homeomorphic to an $(n-1)$ -dimensional ball), let P' be the $(n-1)$ -dimensional regular CW-sphere which is obtained from P by adding an $(n-1)$ -dimensional cell τ so that $\partial\tau = \partial P$.

Definition 2.1. Let P be an $(n-1)$ -dimensional regular CW-sphere. We say that P is *S^* -shellable* if either $P = \{\emptyset\}$ or there is an order $\sigma_1, \sigma_2, \dots, \sigma_r$ of the facets of P such that the following conditions hold.

- (a) $\partial\bar{\sigma}_1$ is S^* -shellable.
- (b) For $1 \leq i \leq r-1$, let

$$\Omega_i = \bar{\sigma}_1 \cup \bar{\sigma}_2 \cup \dots \cup \bar{\sigma}_i$$

and for $2 \leq i \leq r-1$ let

$$\Gamma_i = \overline{[\partial\bar{\sigma}_i \setminus (\partial\bar{\sigma}_i \cap \Omega_{i-1})]}.$$

Then both Ω_i and Γ_i are regular CW-balls of dimension $(n-1)$ and $(n-2)$ respectively, and Γ'_i is S^* -shellable with the first facet of the shelling being the facet which is not in Γ_i .

Remark 2.2. The difference between the above definition and Stanley's S -shellability is that S -shellability only assume that P and Γ'_i are Eulerian and assume no conditions on Ω_i . However, S^* -shellable regular CW-spheres are S -shellable, and the boundary complex of convex polytopes are S^* -shellable by the line shelling [BM]. We leave the verification of this fact to the readers.

The next recursive formula is due to Stanley [St1].

Lemma 2.3 (Stanley). *With the same notation as in Definition 2.1, for $i = 1, 2, \dots, r-2$, one has*

$$\Phi_{\Omega'_{i+1}}(\mathbf{c}, \mathbf{d}) = \Phi_{\Omega'_i}(\mathbf{c}, \mathbf{d}) + \left\{ \Phi_{\Gamma'_{i+1}}(\mathbf{c}, \mathbf{d}) - \Phi_{\Sigma(\partial\Gamma_{i+1})}(\mathbf{c}, \mathbf{d}) \right\} \mathbf{c} + \Phi_{\partial\Gamma_{i+1}}(\mathbf{c}, \mathbf{d}) \mathbf{d}.$$

Since $\Omega'_{r-1} = P$ the above formula gives a way to compute the \mathbf{cd} -index of P recursively.

Next, we recall a result of Ehrenborg and Karu proving that the \mathbf{cd} -index increases by taking subdivisions. Let P and Q be regular CW-complexes, and let $\phi : \mathcal{F}(P) \rightarrow \mathcal{F}(Q)$ be a poset map. For a subcomplex $Q' = \sigma_1 \cup \dots \cup \sigma_s \subset Q$, where each σ_i is a cell of Q , we write $\phi^{-1}(Q') = \phi^{-1}(\sigma_1) \cup \dots \cup \phi^{-1}(\sigma_s)$.

Following [EK, Definition 2.6], for $(n-1)$ -dimensional regular CW-spheres P and \hat{P} , we say that \hat{P} is a subdivision of P if there is an order preserving surjective poset map $\phi : \mathcal{F}(\hat{P}) \rightarrow \mathcal{F}(P)$, satisfying that for any cell σ of P , $\phi^{-1}(\bar{\sigma})$ is a homology ball having the same dimension as σ and $\phi^{-1}(\partial\bar{\sigma}) = \partial(\phi^{-1}(\bar{\sigma}))$.

The following result was proved in [EK, Theorem 1.5].

Lemma 2.4 (Ehrenborg-Karu). *Let P and \hat{P} be $(n-1)$ -dimensional regular CW-spheres. If \hat{P} is a subdivision of P then one has a coefficientwise inequality $\Phi_{\hat{P}}(\mathbf{c}, \mathbf{d}) \geq \Phi_P(\mathbf{c}, \mathbf{d})$*

Back to S^* -shellable regular CW-spheres, with the same notation as in Definition 2.1, Ω'_i is a subdivision of $\Sigma(\partial\Omega_i)$ and $\partial\Omega_i$ is a subdivision of $\Sigma(\partial\Gamma_{i+1})$. Indeed, for the first statement, if τ_1 and τ_2 are the facets of $\Sigma(\partial\Omega_i)$ then define $\phi : \mathcal{F}(\Omega'_i) \rightarrow \mathcal{F}(\Sigma(\partial\Omega_i))$ by

$$\phi(\sigma) = \begin{cases} \sigma, & \text{if } \sigma \in \partial\Omega_i, \\ \tau_1, & \text{if } \sigma \text{ is an interior face of } \Omega_i, \\ \tau_2, & \text{if } \sigma \notin \Omega_i. \end{cases}$$

Similarly, for the second statement, if τ_1 and τ_2 are the facets of $\Sigma(\partial\Gamma_{i+1})$ then define $\phi : \mathcal{F}(\partial\Omega_i) \rightarrow \mathcal{F}(\Sigma(\partial\Gamma_{i+1}))$ by

$$\phi(\sigma) = \begin{cases} \sigma, & \text{if } \sigma \in \partial\Gamma_{i+1}, \\ \tau_1, & \text{if } \sigma \in \bar{\sigma}_{i+1} \setminus \partial\Gamma_{i+1}, \\ \tau_2, & \text{otherwise.} \end{cases}$$

Since $\Phi_{\Sigma P}(\mathbf{c}, \mathbf{d}) = \Phi_P(\mathbf{c}, \mathbf{d})\mathbf{c}$ for any regular CW-sphere P (see [St1, Lemma 1.1]), Lemma 2.4 shows

Lemma 2.5. *With the same notation as in Definition 2.1, for $i = 2, 3, \dots, r-2$, one has $\Phi_{\Omega'_i}(\mathbf{c}, \mathbf{d}) \geq \Phi_{\partial\Gamma_{i+1}}(\mathbf{c}, \mathbf{d})\mathbf{c}^2$.*

3. PROOF OF THE MAIN THEOREM

In this section, we prove Theorem 1.1.

For a homogeneous \mathbf{cd} -polynomial Φ (i.e., homogeneous polynomial of $\mathbb{Z}\langle \mathbf{c}, \mathbf{d} \rangle$ with $\deg \mathbf{c} = 1$ and $\deg \mathbf{d} = 2$) of degree n , we define $\Phi_0, \Phi_2, \dots, \Phi_n$ by

$$\Phi = \Phi_0 + \Phi_2 \mathbf{dc}^{n-2} + \Phi_3 \mathbf{dc}^{n-3} + \dots + \Phi_{n-1} \mathbf{dc} + \Phi_n \mathbf{d}$$

where $\Phi_0 = \alpha \mathbf{c}^n$ for some $\alpha \in \mathbb{Z}$ and each Φ_k is a \mathbf{cd} -polynomial of degree $k - 2$ for $k \geq 2$. Also, we write $\Phi_{\leq k} = \Phi_0 + \Phi_2 \mathbf{dc}^{n-2} + \cdots + \Phi_k \mathbf{dc}^{n-k}$.

Definition 3.1.

- A vector $(\delta_0, \delta_1, \dots, \delta_s) \in \mathbb{Z}^{s+1}$ is said to be k -FFK if there is a k -colored simplicial complex Δ such that $\delta_i = f_{i-1}(\Delta)$ for $i = 0, 1, \dots, s$. ($\{\emptyset\}$ is a 0-colored simplicial complex.) A homogeneous \mathbf{cd} -polynomial $\Phi = \Phi(\mathbf{c}, \mathbf{d})$ is said to be k -FFK if, when we write $\Phi(1, \mathbf{d}) = \delta_0 + \delta_1 \mathbf{d} + \cdots + \delta_s \mathbf{d}^s$, the vector $(\delta_0, \delta_1, \dots, \delta_s)$ is k -FFK.
- A homogeneous \mathbf{cd} -polynomial Φ of degree n is said to be *primitive* if the coefficient of \mathbf{c}^n in Φ is 1.
- Let Φ be a homogeneous \mathbf{cd} -polynomial. A primitive homogeneous \mathbf{cd} -polynomial Ψ is said to be k -good for Φ if Ψ is k -FFK and $\Phi(1, \mathbf{d}) \geq \Psi(1, \mathbf{d})$. Also, we say that a homogeneous \mathbf{cd} -polynomial Ψ is k -good for Φ if it is the sum of primitive homogeneous \mathbf{cd} -polynomials that are k -good for Φ .

We will use the following observation, which follows from [NPT, Lemma 3.1]:

Lemma 3.2. *If Φ is a k -FFK homogeneous \mathbf{cd} -polynomial of degree n , and if Ψ' and Ψ'' are homogeneous \mathbf{cd} -polynomials of degree n' and n'' respectively, where $n', n'' \leq n - 2$, which are k -good for Φ then*

$$\Phi + \Psi' \mathbf{dc}^{n-n'-2} \text{ and } \Phi + \Psi' \mathbf{dc}^{n-n'-2} + \Psi'' \mathbf{dc}^{n-n''-2}$$

are $(k + 1)$ -FFK.

Proof. By Frankl-Füredi-Kalai theorem [FFK], for any k -colored simplicial complex Γ , there is the unique k -colored simplicial complex $\mathcal{C}(\Gamma)$, called a k -colored compressed complex, such that $f_i(\Gamma) = f_i(\mathcal{C}(\Gamma))$ for all i . Moreover, if Γ' is a k -colored complex satisfying $f_i(\Gamma) \leq f_i(\Gamma')$ for all i , then one has $\mathcal{C}(\Gamma) \subset \mathcal{C}(\Gamma')$.

For a simplicial complex Δ , we write $f(\Delta, \mathbf{d}) = 1 + f_0(\Delta) \mathbf{d} + f_1(\Delta) \mathbf{d}^2 + \cdots$. There are k -colored complexes $\Delta, \Delta^{(1)}, \dots, \Delta^{(m)}, \dots, \Delta^{(s)}$ such that $f(\Delta, \mathbf{d}) = \Phi(1, \mathbf{d})$, $\sum_{1 \leq i \leq m} f(\Delta^{(i)}, \mathbf{d}) = \Psi'(1, \mathbf{d})$, $\sum_{m+1 \leq i \leq s} f(\Delta^{(i)}, \mathbf{d}) = \Psi''(1, \mathbf{d})$ and each $\Delta^{(i)}$ is a subcomplex of Δ . Let

$$\Gamma^{(i)} = \Delta \cup \left\{ \bigcup_{k=1}^i \{F \cup \{v_k\} : F \in \Delta^{(k)}\} \right\},$$

where v_1, \dots, v_s are new vertices. Since each $\Delta^{(k)}$ is a subcomplex of Δ , $\Gamma^{(i)}$ is a simplicial complex. Also, $f(\Gamma^{(m)}, \mathbf{d}) = (\Phi + \Psi' \mathbf{dc}^{n-n'-2})(1, \mathbf{d})$ and $f(\Gamma^{(s)}, \mathbf{d}) = (\Phi + \Psi' \mathbf{dc}^{n-n'-2} + \Psi'' \mathbf{dc}^{n-n''-2})(1, \mathbf{d})$. We claim that each $\Gamma^{(i)}$ is $(k + 1)$ -colored. Let V be the vertex set of Δ and $c : V \rightarrow [k]$ a k -coloring map of Δ . Then the map $\hat{c} : V \cup \{v_1, \dots, v_i\} \rightarrow [k + 1]$ defined by $\hat{c}(x) = c(x)$ if $x \in V$ and $\hat{c}(x) = k + 1$ if $x \notin V$ is a $(k + 1)$ -coloring map of $\Gamma^{(i)}$. \square

Let P be an $(n - 1)$ -dimensional S^* -shellable regular CW-sphere with the shelling $\sigma_1, \dots, \sigma_r$. Keeping the notation in Definition 2.1, to simplify notations, we use the

following symbols.

$$\begin{aligned}\Phi^{(i)} &= \Phi^{(i)}(\mathbf{c}, \mathbf{d}) = \Phi_{\Omega'_i}(\mathbf{c}, \mathbf{d}) \\ \Phi &= \Phi_P(\mathbf{c}, \mathbf{d}) = \Phi^{(r-1)} \\ \Psi^{(i)} &= \Phi_{\Gamma'_{i+1}}(\mathbf{c}, \mathbf{d}) - \Phi_{\Sigma(\partial\Gamma_{i+1})}(\mathbf{c}, \mathbf{d}) \\ \Psi &= \sum_{i=1}^{r-2} \Psi^{(i)} \\ \Pi &= \Phi - \Phi^{(1)}.\end{aligned}$$

Thus Stanley's recursive formula, Lemma 2.3, says

$$\Phi^{(i+1)} = \Phi^{(i)} + \Psi^{(i)}\mathbf{c} + \Phi_{\partial\Gamma_{i+1}}\mathbf{d}$$

and

$$\Pi = \Psi\mathbf{c} + \sum_{i=1}^{r-2} \Phi_{\partial\Gamma_{i+1}}(\mathbf{c}, \mathbf{d})\mathbf{d}.$$

The last part of the following proposition is a restatement of Theorem 1.1.

Proposition 3.3. *With notation as above, the following holds.*

- (1) For $2 \leq k \leq n$, $\Psi_k^{(i)}$ is $\lfloor \frac{k}{2} - 1 \rfloor$ -good for $\Phi_{\leq k-2}^{(i)} + \Psi_{\leq k-2}^{(i)}\mathbf{c}$.
- (2) For $2 \leq k \leq n$, Π_k is $\lfloor \frac{k}{2} - 1 \rfloor$ -good for $\Phi_{\leq k-2}^{(1)} + \Pi_{\leq k-2}$.
- (3) For $2 \leq k \leq n$, Φ_k is $\lfloor \frac{k}{2} - 1 \rfloor$ -good for $\Phi_{\leq k-2}$.
- (4) For $0 \leq k \leq n$, $\Phi_{\leq k}$ is $\lfloor \frac{k}{2} \rfloor$ -FFK. In particular, the \mathbf{cd} -index of P is $\lfloor \frac{n}{2} \rfloor$ -FFK.

Proof. The proof is by induction on dimension, where all statements clearly hold for $n = 0, 1$. Suppose that all statements are true up to dimension $n - 2$. To simplify notations, for a regular CW-sphere Q , we write $\Phi_Q = \Phi_Q(\mathbf{c}, \mathbf{d})$.

Proof of (1). By applying the induction hypothesis to Γ'_{i+1} (use statement(2)), each $\Psi_k^{(i)}$ is $\lfloor \frac{k}{2} - 1 \rfloor$ -good for $(\Phi_{\Sigma(\partial\Gamma_{i+1})})_{\leq k-2}^{(i)} + \Psi_{\leq k-2}^{(i)}$. Since $(\Upsilon\mathbf{c})_k = \Upsilon_k$ for any homogeneous \mathbf{cd} -polynomial Υ , $\Psi_k^{(i)}$ is $\lfloor \frac{k}{2} - 1 \rfloor$ -good for $(\Phi_{\Sigma(\partial\Gamma_{i+1})})_{\leq k-2}^{(i)}\mathbf{c} + \Psi_{\leq k-2}^{(i)}\mathbf{c}$. By Lemma 2.5,

$$\Phi_{\Sigma(\partial\Gamma_{i+1})}\mathbf{c} = \Phi_{\partial\Gamma_{i+1}}\mathbf{c}^2 \leq \Phi_{\Omega'_i} = \Phi^{(i)},$$

thus $\Psi_k^{(i)}$ is $\lfloor \frac{k}{2} - 1 \rfloor$ -good for $\Phi_{\leq k-2}^{(i)} + \Psi_{\leq k-2}^{(i)}\mathbf{c}$.

Proof of (2). By the definition of Π ,

$$\Pi_k = \sum_{i=1}^{r-1} \Psi_k^{(i)} \text{ for } k < n$$

and

$$\Pi_n = \sum_{i=1}^{r-2} \Phi_{\partial\Gamma_{i+1}}.$$

By (1), each $\Psi_k^{(i)}$ is $\lfloor \frac{k}{2} - 1 \rfloor$ -good for $\Phi_{\leq k-2}^{(i)} + \Psi_{\leq k-2}^{(i)}\mathbf{c}$. Then since

$$\Phi_{\leq k-2}^{(i)} + \Psi_{\leq k-2}^{(i)}\mathbf{c} \leq \Phi_{\leq k-2} = \Phi_{\leq k-2}^{(1)} + \Pi_{\leq k-2},$$

Π_k is $\lfloor \frac{k}{2} - 1 \rfloor$ -good for $\Phi_{\leq k-2}^{(1)} + \Pi_{\leq k-2}$ for $k < n$. Also, each $\Phi_{\partial\Gamma_{i+1}}$ is $\lfloor \frac{n}{2} - 1 \rfloor$ -FFK by the induction hypothesis (use (4)), and $\Phi_{\partial\Gamma_{i+1}} \mathbf{c}^2 \leq \Phi^{(i)}$ by Lemma 2.5. The latter condition clearly says

$$\Phi_{\partial\Gamma_{i+1}} \mathbf{c}^2 \leq \Phi_{\leq n-2}^{(i)} \leq \Phi_{\leq n-2} = \Phi_{\leq n-2}^{(1)} + \Pi_{\leq n-2}.$$

Hence Π_n is $\lfloor \frac{n}{2} - 1 \rfloor$ -good for $\Phi_{\leq n-2}^{(1)} + \Pi_{\leq n-2}$.

Proof of (3). Observe that since $\Phi^{(1)} = \Phi_{\partial\bar{\sigma}_1} \mathbf{c}$,

$$\Phi_k = \Phi_k^{(1)} + \Psi_k \text{ for } k < n$$

and

$$\Phi_n = \Pi_n.$$

We already proved that $\Phi_n = \Pi_n$ is $\lfloor \frac{n}{2} - 1 \rfloor$ -good for $\Phi_{\leq n-2}$ in the proof of (2). Suppose $k < n$. Since $\Phi^{(1)} = \Phi_{\partial\bar{\sigma}_1} \mathbf{c}$, by the induction hypothesis (use (3)), $\Phi_k^{(1)}$ is $\lfloor \frac{k}{2} - 1 \rfloor$ -good for $\Phi_{\leq k-2}^{(1)}$. Since $\Phi_{\leq k-2}^{(1)} \leq \Phi_{\leq k-2}$ and since we already proved that $\Psi_k = \Pi_k$ is $\lfloor \frac{k}{2} - 1 \rfloor$ -good for $\Phi_{\leq k-2}$ in the proof of (2), Φ_k is $\lfloor \frac{k}{2} - 1 \rfloor$ -good for $\Phi_{\leq k-2}$.

Proof of (4). This statement easily follows from (3). For $k = 0, 1$, the statement is obvious (as $\Phi_{\leq 0} = \Phi_{\leq 1} = \mathbf{c}^n$). Suppose that $\Phi_{\leq 2m+1}$ is m -FFK, where $m \in \mathbb{Z}_{\geq 0}$. Then both Φ_{2m+2} and Φ_{2m+3} are m -good for $\Phi_{\leq 2m+1}$ by (3), and therefore $\Phi_{\leq 2m+2}$ and $\Phi_{\leq 2m+3}$ are $(m+1)$ -FFK by Lemma 3.2. \square

4. γ -VECTORS OF POLYTOPES AND A CONJECTURE ON THE \mathbf{cd} -INDEX

γ -vectors and the \mathbf{cd} -index. Let Δ be an $(n-1)$ -dimensional simplicial complex. Then the h -vector $h(\Delta) = (h_0, h_1, \dots, h_n)$ of Δ is defined by the relation

$$\sum_{i=0}^n h_i x^{n-i} = \sum_{i=0}^n f_{i-1}(\Delta) (x-1)^{n-i}.$$

If Δ is a simplicial sphere (that is, a triangulation of a sphere), or more generally a homology sphere, then $h_i = h_{n-i}$ for all i by the Dehn-Sommerville equations, and in this case the γ -vector $(\gamma_0, \gamma_1, \dots, \gamma_{\lfloor \frac{n}{2} \rfloor})$ of Δ is defined by the relation

$$\sum_{i=0}^n h_i x^i = \sum_{i=0}^{\lfloor \frac{n}{2} \rfloor} \gamma_i x^i (1+x)^{n-2i}.$$

It was conjectured by Gal [Ga] that if Δ is a flag homology sphere then its γ -vector is non-negative. Recently Nevo and Peterson [NP] further conjectured that the γ -vector of a flag homology sphere is the f -vector of a balanced simplicial complex. These conjectures are open in general, the latter conjecture was verified for barycentric subdivisions of simplicial homology spheres [NPT], and Gal's conjecture is known to be true for barycentric subdivisions of regular CW-spheres by the following fact, combined with Karu's result on the nonnegativity of the \mathbf{cd} -index for Gorenstien* posets:

Let P be an $(n-1)$ -dimensional regular CW-sphere. The *barycentric subdivision* $\text{sd}(P)$ of P is the order complex of $\mathcal{F}(P)$. Let (h_0, h_1, \dots, h_n) and $(\gamma_0, \gamma_1, \dots, \gamma_{\lfloor \frac{n}{2} \rfloor})$ be the h -vector and γ -vector of $\text{sd}(P)$, respectively. Then it is easy to see that

$h_i = \sum_{S \subset [n], |S|=i} h_S(P)$. Thus if $\Phi_P(1, \mathbf{d}) = \delta_0 + \delta_1 \mathbf{d} + \delta_2 \mathbf{d}^2 + \cdots + \delta_{\lfloor \frac{n}{2} \rfloor} \mathbf{d}^{\lfloor \frac{n}{2} \rfloor}$, then for all $i \geq 0$,

$$\gamma_i = 2^i \delta_i.$$

Since δ_i is non-negative, we conclude that γ_i is also non-negative.

The next simple statement, combined with Theorem 1.1, proves Theorem 1.2.

Lemma 4.1. *With the same notation as above, if $(\delta_0, \delta_1, \dots, \delta_{\lfloor \frac{n}{2} \rfloor})$ is k -FFK then $(\gamma_0, \gamma_1, \dots, \gamma_{\lfloor \frac{n}{2} \rfloor})$ is also k -FFK.*

Proof. Let Δ be a k -colored simplicial complex on the vertex set V with $f_{i-1}(\Delta) = \delta_i$ for all $i \geq 0$ and let $c : V \rightarrow [k]$ be a k -coloring map of Δ . Consider a collection of subsets of $W = \{x_v : v \in V\} \cup \{y_v : v \in V\}$

$$\hat{\Delta} = \{x_G \cup y_{F \setminus G} : F \in \Delta, G \subset F\},$$

where $x_H = \{x_v : v \in H\}$ and $y_H = \{y_v : v \in H\}$ for any $H \subset V$. Then $\hat{\Delta}$ is a simplicial complex with $f_{i-1}(\hat{\Delta}) = 2^i f_{i-1}(\Delta) = \gamma_i$ for all i . The map $\hat{c} : W \rightarrow [k]$, $\hat{c}(x_v) = \hat{c}(y_v) = c(v)$, shows that $\hat{\Delta}$ is k -colored. \square

Proof of Corollary 1.3. By Theorem 1.2, in order to prove Corollary 1.3 it is enough to show that $\delta_{\lfloor \frac{n}{2} \rfloor}(P) > 0$ where P is the boundary complex of an n -polytope. Billera and Ehrenborg showed that the \mathbf{cd} -index of n -polytopes is minimized (coefficientwise) by the n -simplex, denoted σ^n [BE]. Thus, it is enough to verify that $\delta_{\lfloor \frac{n}{2} \rfloor}(\sigma^n) > 0$. It is known that *all* the \mathbf{cd} -coefficients of σ^n are positive (e.g., by using the Ehrenborg-Readdy formula for the \mathbf{cd} -index of a pyramid over a polytope [ER, Theorem 5.2]). \square

A conjecture on the \mathbf{cd} -index. It would be natural to ask if Theorems 1.1 and 1.2 hold for all regular CW-spheres (or all Gorenstein* posets). We phrase a conjecture on the the \mathbf{cd} -index, that, if true, immediately implies Theorem 1.1, as well as the entire Proposition 3.3(4).

For an arbitrary \mathbf{cd} -monomial $w = \mathbf{c}^{s_0} \mathbf{d} \mathbf{c}^{s_1} \mathbf{d} \cdots \mathbf{d} \mathbf{c}^{s_k}$ of degree n (where $0 \leq s_i$ for all i and $s_0 + \cdots + s_k + 2k = n$), let F_w be the following subset of $[n-1]$:

$$F_w = \{s_0 + 1, s_0 + s_1 + 3, s_0 + s_1 + s_2 + 5, \dots, s_0 + \cdots + s_{k-1} + 2k - 1\}.$$

Note that F_w contains no two consecutive numbers. For example, $F_{\mathbf{c}^n} = \emptyset$, $F_{\mathbf{d}^k} = \{1, 3, \dots, 2k-1\}$ and $F_{\mathbf{cd}^k} = \{2, 4, \dots, 2k\}$. Let \mathcal{A} be the set of subsets of $[n-1]$ that have no two consecutive numbers, and let \mathcal{B} be the set of \mathbf{cd} -monomials of degree n . Then $w \mapsto F_w$ is a bijection from \mathcal{B} to \mathcal{A} (as $k = |F_w|$ and $s_k = n - 2k - s_{k-1} - \cdots - s_0$ we see that the inverse map exists).

Let Δ be a k -colored simplicial complex with the vertex set V and a k -coloring map $c : V \rightarrow [k]$. For any subset $S \subset [k]$, let $f_S(\Delta) = |\{F \in \Delta : c(F) = S\}|$. The vector $(f_S(\Delta) : S \subset [k])$ is called the *flag f -vector* of Δ . Note that the flag f -vector of a Gorenstein* poset P is equal to the flag f -vector of $\text{sd}(P)$ by the coloring map defined by the rank function.

Definition 4.2. Let $\Phi = \sum_w a_w w$ be a homogeneous \mathbf{cd} -polynomial of degree n with w the \mathbf{cd} -monomials and $a_w \in \mathbb{Z}$. For $S \subset [n-1]$, we define

$$\alpha_S(\Phi) = \begin{cases} a_w, & \text{if } S = F_w \text{ for some } w \in \mathcal{B} \\ 0, & \text{if } S \notin \mathcal{A}. \end{cases}$$

Conjecture 4.3. Let P be an $(n - 1)$ -dimensional regular CW-sphere (or more generally, Gorenstein* poset of rank $n + 1$). Then there exists an $(n - 1)$ -colored simplicial complex Δ such that $f_S(\Delta) = \alpha_S(\Phi_P)$ for all $S \subset [n - 1]$.

Thus the above conjecture states that the \mathbf{cd} -index is itself the flag f -vector of a colored complex. If the above conjecture is true then $\Phi_P(1, \mathbf{d}) = 1 + f_0(\Delta)\mathbf{d} + \cdots + f_{\lfloor \frac{n}{2} \rfloor - 1}(\Delta)\mathbf{d}^{\lfloor \frac{n}{2} \rfloor}$. Although Δ is $(n - 1)$ -colored, this fact implies Theorem 1.1. Indeed, since $f_S(\Delta) = \alpha_S(\Phi_P) = 0$ if S has consecutive numbers, if $c : V \rightarrow [n - 1]$ is an $(n - 1)$ -coloring map of Δ then the map $\hat{c} : V \rightarrow [\lfloor \frac{n}{2} \rfloor]$ defined by $\hat{c}(v) = \lfloor \frac{c(v)+1}{2} \rfloor$ is an $\lfloor \frac{n}{2} \rfloor$ -coloring map of Δ .

The next result supports the conjecture in low dimension.

Proposition 4.4. Let P be a Gorenstein* poset of rank $n + 1$. For all $i, j \in [n - 1]$,

$$\alpha_{\{i\}}(\Phi_P)\alpha_{\{j\}}(\Phi_P) \geq \alpha_{\{i,j\}}(\Phi_P).$$

Proof. Let $(h_S(P) : S \subset [n])$ be the flag h -vector of P . Let $\{i, i + j\} \subset [n - 1]$ with $j \geq 2$. What we must prove is $\alpha_{\{i\}}(\Phi_P)\alpha_{\{i+j\}}(\Phi_P) \geq \alpha_{\{i,i+j\}}(\Phi_P)$.

Observe that

$$\begin{aligned} h_{[i] \cup \{i+j+1, \dots, n\}}(P) &= \alpha_{\{i,i+j\}}(\Phi_P) + \alpha_{\{i\}}(\Phi_P) + \alpha_{\{i+j\}}(\Phi_P) + \alpha_\emptyset(\Phi_P), \\ h_{[i]}(P) &= \alpha_{\{i\}}(\Phi_P) + \alpha_\emptyset(\Phi_P), \\ h_{\{i+j+1, \dots, n\}}(P) &= \alpha_{\{i+j\}}(\Phi_P) + \alpha_\emptyset(\Phi_P) \end{aligned}$$

(as $h_{[i] \cup \{i+j+1, \dots, n\}}(P)$ is the coefficient of $\mathbf{b}^i \mathbf{a}^j \mathbf{b}^{n-i-j}$ in $\Psi_P(\mathbf{a}, \mathbf{b})$, etc.). Since $\alpha_\emptyset = 1$, it is enough to prove that

$$h_{[i]}(P)h_{\{i+j+1, \dots, n\}}(P) \geq h_{[n-i-j] \cup \{n-i+1, \dots, n\}}(P).$$

It follows from [St2, III, Theorem 4.6] that there is an n -colored simplicial complex Δ with a coloring map $c : V \rightarrow [n]$ such that $f_S(\Delta) = h_S(P)$ for all $S \subset [n]$. Let

$$\Delta_S = \{F \in \Delta : c(F) = S\}$$

for $S \subset [n]$. Then it is clear that

$$\Delta_{[i] \cup \{i+j+1, \dots, n\}} \subset \{F \cup G : F \in \Delta_{[i]}, G \in \Delta_{\{i+j+1, \dots, n\}}\},$$

which implies the desired inequality. \square

It is straightforward that the above proposition proves the next statement.

Corollary 4.5. Conjecture 4.3 holds for $n \leq 5$.

Non-existence of \mathbf{d} -polynomials. For a Gorenstein* poset P , we call $\Phi_P(1, \mathbf{d})$ the \mathbf{d} -polynomial of P . It is a challenging problem to classify all possible \mathbf{d} -polynomials of Gorenstein* posets, which give a complete characterization of all possible face vectors of Gorenstein* order complexes since knowing \mathbf{d} -polynomials is equivalent to knowing γ -vectors. The problem is open even for the 3-dimensional case. To study this problem, by virtue of Theorem 1.1, it is natural to ask which FFK vector is realizable as the \mathbf{d} -polynomial of a Gorenstein* poset. The next result shows that not all $\lfloor \frac{n}{2} \rfloor$ -FFK vectors are realizable as the \mathbf{d} -polynomial of a Gorenstein* poset of rank $n + 1$.

First recall that the ordinal sum $Q_1 + Q_2$ of two disjoint posets Q_1 and Q_2 is the poset whose elements are the union of elements in Q_1 and Q_2 and whose relations

are those in Q_1 union those in Q_2 union all $q_1 < q_2$ where $q_1 \in Q_1$ and $q_2 \in Q_2$. For Gorenstein* posets Q_1 and Q_2 , the poset $Q_1 * Q_2 = (Q_1 - \{\hat{1}\}) + (Q_2 - \{\hat{0}\})$ is called the *join* of Q_1 and Q_2 , and $\Sigma Q_1 = Q_1 * B_2$, where B_2 is a Boolean algebra of rank 2, is called the *suspension* of Q_1 . By [St1, Lemma 1.1], $\Phi_{Q_1 * Q_2}(\mathbf{c}, \mathbf{d}) = \Phi_{Q_1}(\mathbf{c}, \mathbf{d}) \cdot \Phi_{Q_2}(\mathbf{c}, \mathbf{d})$.

Proposition 4.6. *Let P be a Gorenstein* poset of rank 5, and let*

$$\Phi_P(\mathbf{c}, \mathbf{d}) = \mathbf{c}^4 + \alpha_{\{1\}} \mathbf{c}^2 \mathbf{d} + \alpha_{\{2\}} \mathbf{c} \mathbf{d} \mathbf{c} + \alpha_{\{3\}} \mathbf{d} \mathbf{c}^2 + \alpha_{\{1,3\}} \mathbf{d}^2$$

be its \mathbf{cd} -index. Suppose $\alpha_{\{2\}} = 0$. Then there are Gorenstein posets P_1 and P_2 of rank 3 such that $P = P_1 * P_2$. In particular, $\alpha_{\{1,3\}} = \alpha_{\{1\}} \alpha_{\{3\}}$.*

Proof. Let r denote the rank function $r : P \rightarrow \{0, 1, \dots, 5\}$ ($r(\hat{0}) = 0, r(\hat{1}) = 5$). Let $P_1 := \{F \in P : r(F) \leq 2\}$ and $P_2 := \{F \in P : r(F) \geq 3\}$.

As P is Gorenstien*, to show that $P = P_1 + P_2$ it is enough to show that $P_2 \cup \{\hat{0}\}$ is Gorenstien* (as a Gorenstien* poset contains no proper subposet which is Gorenstien* of the same rank, and each interval $[F, \hat{1}]$ with $r(F) = 2$ in P is Gorenstien*). For this, it is enough to show that any rank 4 element in P covers exactly two rank 3 elements in P . Indeed, this guarantees that the dual poset to P_2 , denoted P_2^* , is the face poset of a union of CW 1-spheres, and as P is Gorenstien* so is its dual P^* , hence P_2^* is Cohen-Macaulay since P_2^* is a rank selected poset [St2, III, Theorem 4.5], which implies that P_2^* is the face poset of one CW 1-sphere, i.e. $P_2 \cup \{\hat{0}\}$ is Gorenstien*.

Let F be a rank 4 element of P . Then P is a subdivision of $\Sigma([\hat{0}, F])$ (Recalling [EK, Definition 2.6], this is shown by the map $\phi : P \rightarrow \Sigma([\hat{0}, F])$, $\phi(\sigma) = \sigma$ if $\sigma < F$, $\phi(\sigma) = \sigma_1$ if σ and F are incomparable, and $\phi(F) = \sigma_2$, where σ_1, σ_2 are the rank 4 elements in $\Sigma([\hat{0}, F])$). Thus, by Lemma 2.4, the coefficient of \mathbf{cdc} in the \mathbf{cd} -index of $\Sigma([\hat{0}, F])$ is zero, hence the coefficient of the monomial \mathbf{cd} in the \mathbf{cd} -index of $[\hat{0}, F]$ is zero.

This fact implies, when expanding the \mathbf{cd} -index of $[\hat{0}, F]$ in terms of \mathbf{a}, \mathbf{b} , that $h_{\{3\}}([\hat{0}, F])$ equals the coefficient of \mathbf{c}^3 , namely $h_{\{3\}}([\hat{0}, F]) = 1$. Switching to the flag f -vector of $[\hat{0}, F]$ we get $f_{\{3\}}([\hat{0}, F]) = h_{\emptyset}([\hat{0}, F]) + h_{\{3\}}([\hat{0}, F]) = 1 + 1 = 2$. Thus, F covers exactly two rank 3 elements in P . \square

Example 4.7. Consider the 2-FFK vector $(1, 6, 7)$. We claim that $\Phi_P(1, \mathbf{d}) \neq 1 + 6\mathbf{d} + 7\mathbf{d}^2$ for all Gorenstein* poset P of rank 5. Indeed, if $\Phi_P(1, \mathbf{d}) = 1 + 6\mathbf{d} + 7\mathbf{d}^2$, then $\alpha_{\{1,3\}} = 7$. Then $\alpha_{\{1\}} + \alpha_{\{3\}} = 6$ and $\alpha_{\{2\}} = 0$ by Proposition 4.4, which contradicts Proposition 4.6.

A similar argument shows that $(1, 2a, a^2 - 2)$, where $a \geq 3$, is 2-FFK, but not realizable as the \mathbf{d} -polynomial of a Gorenstein* poset of rank 5.

REFERENCES

- [BK] M.M. Bayer and A. Klapper, A new index for polytopes, *Discrete Comput. Geom.* **6** (1991), 33–47.
- [BE] L. J. Billera and R. Ehrenborg, Monotonicity of the \mathbf{cd} -index for polytopes, *Math. Z.* **233** (2000), 421–441.
- [BM] H. Bruggesser and P. Mani, Shellable decompositions of cells and spheres, *Math. Scand.* **29** (1971), 197–205.

- [EK] R. Ehrenborg and K. Karu, Decomposition theorem for the cd -index of Gorenstein posets, *J. Algebraic Combin.* **26** (2007), 225–251.
- [ER] R. Ehrenborg and M. Readdy, Coproducts and the cd -index, *J. Algebraic Combin.* **8** (1998), 273–299.
- [FFK] P. Frankl, Z. Füredi and G. Kalai, Shadows of colored complexes, *Math. Scand.* **63** (1998), 169–178.
- [Ga] Š.R. Gal, Real root conjecture fails for five- and higher-dimensional spheres, *Discrete Comput. Geom.* **34** (2005), 269–284.
- [Ka] K. Karu, The cd -index of fans and posets, *Compos. Math.* **142** (2006), 701–718.
- [NP] E. Nevo and T.K. Petersen, On γ -vectors satisfying the Kruskal-Katona inequalities, *Discrete Comput. Geom.*, to appear.
- [NPT] E. Nevo, K.T. Petersen and B. Tenner, The γ -vector of a barycentric subdivision, to appear in *J. Combin. Theory Ser. A*, arXiv:1003.2544q (2010).
- [St1] R.P. Stanley. Flag f -vectors and the cd -index, *Math. Z.* **216** (1994), 483–499.
- [St2] R.P. Stanley, Combinatorics and commutative algebra, Second edition, Progr. Math., vol. 41, Birkhäuser, Boston, 1996.

SATOSHI MURAI, DEPARTMENT OF MATHEMATICAL SCIENCE, FACULTY OF SCIENCE, YAMAGUCHI UNIVERSITY, 1677-1 YOSHIDA, YAMAGUCHI 753-8512, JAPAN

DEPARTMENT OF MATHEMATICS, BEN GURION UNIVERSITY OF THE NEGEV, BE'ER SHEVA 84105, ISRAEL

E-mail address: `nevoe@math.bgu.ac.il`