

# On Embeddability and Stresses of Graphs

Eran Nevo \*

January 2, 2007

## Abstract

Gluck [6] has proven that triangulated 2-spheres are generically 3-rigid. Equivalently, planar graphs are generically 3-stress free. We show that already the  $K_5$ -minor freeness guarantees the stress freeness. More generally, we prove that every  $K_{r+2}$ -minor free graph is generically  $r$ -stress free for  $1 \leq r \leq 4$ . (This assertion is false for  $r \geq 6$ .) Some further extensions are discussed.

## 1 Introduction and results

Gluck [6] has proven that triangulated 2-spheres are generically 3-rigid. His proof is based on two classical theorems: one is Cauchy's rigidity theorem [3] (for convex 3-polytopes), the other is Steinitz's theorem [17] which asserts that any polyhedral 2-sphere is combinatorially isomorphic to the boundary complex of some convex 3-polytope. Whiteley [18] has found a proof of Gluck's theorem which avoids convexity, based on vertex splitting. This proof inspired our main result, Theorem 1.2 below.

It is easy to see that a graph with  $n$  vertices and  $3n-6$  edges is generically 3-rigid iff it is generically 3-stress free. Thus, Gluck's theorem can be stated as:

**Theorem 1.1** (*Gluck*) *Planar graphs are generically 3-stress free.*

Bearing in mind Kuratowski's theorem [9], we show that already the  $K_5$ -minor freeness guarantees the 3-stress freeness, and more generally:

**Theorem 1.2** *For  $2 \leq r \leq 6$ , every  $K_r$ -minor free graph is generically  $(r-2)$ -stress free.*

The proof is based on contracting edges which possess a certain property. It makes an essential use of Mader's theorem [12] which gives an upper bound  $(r-2)n - \binom{r-1}{2}$  on the number of edges in a  $K_r$ -minor free graph with  $n$

---

\*Institute of Mathematics, Hebrew University of Jerusalem, Jerusalem Israel, E-mail address: eranevo@math.huji.ac.il. Supported by an I.S.F. grant.

vertices, for  $r \leq 7$ . Indeed, Theorem 1.2 can be regarded as a strengthening of Mader's theorem, as being generically  $l$ -stress free implies having at most  $ln - \binom{l+1}{2}$  edges (as the rank of the corresponding rigidity matrix is at most  $ln - \binom{l+1}{2}$ , see Section 2). This also shows that Theorem 1.2 fails for  $r \geq 8$ , as is demonstrated for  $r = 8$  by  $K_{2,2,2,2,2}$ , and for  $r > 8$  by repeatedly coning over  $K_{2,2,2,2,2}$  (e.g. [16]). It would be interesting to find a proof of Theorem 1.2 that avoids using Mader's theorem (and derive Mader's theorem as a corollary).

**Problem 1.3** *Must a graph with a generic  $(r - 2)$ -stress contain a subdivision of  $K_r$  for  $2 \leq r \leq 6$ ?*

The answer is positive for  $r = 2, 3, 4$  as in this case  $G$  has a  $K_r$  minor iff  $G$  contains a subdivision of  $K_r$  ([5], Proposition 1.7.2). Mader proved that every graph on  $n$  vertices with more than  $3n - 6$  edges contains a subdivision of  $K_5$  [13]. A positive answer in the case  $r = 5$  would strengthen this result.

A graph is *linklessly embeddable* if there exists an embedding of it in  $\mathbb{R}^3$  (where vertices and edges have disjoint images) such that every two disjoint cycles of it are unlinked closed curves in  $\mathbb{R}^3$ . As such graph is  $K_6$ -minor free (e.g. [15], [11]), combining with Theorem 1.2 we conclude:

**Corollary 1.4** *Linklessly embeddable graphs are generically 4-stress free.*

Let  $\mu(G)$  denote the Colin de Verdière's parameter of a graph  $G$  [4]. Colin de Verdière [4] proved that a graph  $G$  is planar iff  $\mu(G) \leq 3$ ; Lovász and Schrijver [11] proved that  $G$  is linklessly embeddable iff  $\mu(G) \leq 4$ . While we have seen that Theorem 1.2 fails for  $r \geq 8$ , we conjecture that Theorem 1.1 and Corollary 1.4 extend to:

**Conjecture 1.5** *Let  $G$  be a graph and let  $k$  be a positive integer. If  $\mu(G) \leq k$  then  $G$  is generically  $k$ -stress free.*

For  $k = 1, 2, 3, 4$  this is true: Colin de Verdière [4] showed that the family  $\{G : \mu(G) \leq k\}$  is closed under taking minors for every  $k$ . Note that  $\mu(K_r) = r - 1$ . By Theorem 1.2 Conjecture 1.5 holds for  $k \leq 4$ . Conjecture 1.5 implies

$$\mu(G) \leq k \Rightarrow e \leq kv - \binom{k+1}{2}$$

(where  $e$  and  $v$  are the numbers of edges and vertices in  $G$ , respectively) which is not known either.

This paper is organized as follows: Section 2 provides relevant background in rigidity theory of graphs, Section 3 deals with graph minors, and in Section 4 we prove Theorem 1.2.

## 2 Rigidity

The presentation here is based mainly on Kalai's [7]. Let  $G = (V, E)$  be a graph. Let  $d(a, b)$  denote Euclidian distance between points  $a$  and  $b$  in Euclidian space. A  $d$ -embedding  $f : V \rightarrow \mathbb{R}^d$  is called *rigid* if there exists an  $\varepsilon > 0$  such that if  $g : V \rightarrow \mathbb{R}^d$  satisfies  $d(f(v), g(v)) < \varepsilon$  for every  $v \in V$  and  $d(g(u), g(w)) = d(f(u), f(w))$  for every  $\{u, w\} \in E$ , then  $d(g(u), g(w)) = d(f(u), f(w))$  for every  $u, w \in V$ . Loosely speaking,  $f$  is rigid if any perturbation of it which preserves the lengths of the edges is induced by an isometry of  $\mathbb{R}^d$ .  $G$  is called *generically  $d$ -rigid* if the set of its rigid  $d$ -embeddings is open and dense in the topological vector space of all of its  $d$ -embeddings. Given a  $d$ -embedding  $f : V \rightarrow \mathbb{R}^d$ , a *stress* w.r.t.  $f$  is a function  $w : E \rightarrow \mathbb{R}$  such that for every vertex  $v \in V$

$$\sum_{w:\{v,u\} \in E} w(\{v,u\})(f(v) - f(u)) = 0.$$

$G$  is called *generically  $d$ -stress free* if the set of its  $d$ -embeddings which have a unique stress ( $w = 0$ ) is open and dense in the space of all of its  $d$ -embeddings.

Rigidity and stress freeness can be related as follows: Let  $V = [n]$ , and let  $Rig(G, f)$  be the  $dn \times |E|$  matrix associated with a  $d$ -embedding  $f$  of  $V(G)$  defined as follows: for its column corresponding to  $\{v < u\} \in E$  put the vector  $f(v) - f(u)$  (resp.  $f(u) - f(v)$ ) at the entries of the rows corresponding to  $v$  (resp.  $u$ ) and zero otherwise.  $G$  is generically  $d$ -stress free if  $Ker(Rig(G, f)) = 0$  for a generic  $f$  (i.e. for an open dense set of embeddings).  $G$  is generically  $d$ -rigid if  $Im(Rig(G, f)) = Im(Rig(K_V, f))$  for a generic  $f$ , where  $K_V$  is the complete graph on  $V = V(G)$ . The dimensions of the kernel and image of  $Rig(G, f)$  are independent of the generic  $f$  we choose; we call  $R(G) = Rig(G, f)$  the *rigidity matrix* of  $G$ . For the complete graph, one computes  $rank(Rig(K_V)) = dn - \binom{d+1}{2}$  (see Asimov and Roth [1] for more details).

We need the following two known results:

**Theorem 2.1** (Asimov and Roth [2]) *Let  $G_i = (V_i, E_i)$  be generically  $k$ -stress free graphs,  $i = 1, 2$  such that  $G_1 \cap G_2$  is generically  $k$ -rigid. Then  $G_1 \cup G_2$  is generically  $k$ -stress free.*

**Theorem 2.2** (Whiteley [18]) *Let  $G'$  be obtained from a graph  $G$  by contracting an edge  $\{u, v\}$ .*

(a) *If  $u, v$  have at least  $d - 1$  common neighbors and  $G'$  is generically  $d$ -rigid, then  $G$  is generically  $d$ -rigid.*

(b) *If  $u, v$  have at most  $d - 1$  common neighbors and  $G'$  is generically  $d$ -stress free, then  $G$  is generically  $d$ -stress free.*

Theorem 2.2 gives an alternative proof of Gluck's theorem (Whiteley [18]): starting with a triangulated 2-sphere, repeatedly contract edges with exactly 2 common neighbors until a tetrahedron is reached (it is not difficult to show that this is always possible). By Theorem 2.2(a) it is enough to show that the tetrahedron is generically 3-rigid, as is well known (Asimov and Roth [1]).

**Remark:** Generic rigidity is equivalent to a certain property of Kalai's symmetric algebraic shifting of the graph, see [8] and [10] for details. Kalai defined exterior shifting as well; see [8] for details and references. Let  $\Delta$  denote the algebraic shifting operator, for both symmetric and exterior versions. Theorem 1.2 extends to:

*for every  $2 \leq r \leq 6$  and every graph  $G$ , if  $\{r-1, r\} \in \Delta(G)$  then  $G$  has a  $K_r$  minor.*

Details can be found in early version of this paper [14]. In relation with Conjecture 1.5, we conjecture  $\mu(G) \leq k \Rightarrow \{k+1, k+2\} \notin \Delta(K)$ . The early version [14] also discusses connections with embeddability into 2-manifolds.

### 3 Minors

All graphs we consider are simple, i.e. with no loops and no multiple edges. Let  $e = \{v, u\}$  be an edge in a graph  $G$ . By *contracting*  $e$  we mean identifying the vertices  $v$  and  $u$  and deleting the loop and one copy of each double edge created by this identification, to obtain a new (simple) graph. A graph  $H$  is called a *minor* of a graph  $G$ , denoted  $H \prec G$ , if by repeated contraction of edges we can obtain  $H$  from a subgraph of  $G$ . In the sequel we shall make an essential use of the following Theorem of Mader:

**Theorem 3.1** (Mader [12]) *For  $3 \leq r \leq 7$ , if a graph  $G$  on  $n$  vertices has no  $K_r$  minor then it has at most  $(r-2)n - \binom{r-1}{2}$  edges.*

**Proposition 3.2** *For  $3 \leq r \leq 5$ : If  $G$  has an edge and each edge belongs to at least  $r-2$  triangles, then  $G$  has a  $K_r$  minor.*

*Proof:* For  $r = 3$   $G$  actually contains  $K_3$  as a subgraph. Let  $G$  have  $n$  vertices and  $e$  edges. Assume (by contradiction) that  $K_r \not\prec G$ . W.l.o.g.  $G$  is connected.

For  $r = 4$ , by Theorem 3.1  $e \leq 2n - 3$  hence there is a vertex  $u \in G$  with degree  $d(u) \leq 3$ . Denote by  $N(u)$  the induced subgraph on the neighbors of  $u$ . For every  $v \in N(u)$ , the edge  $uv$  belongs to at least two triangles, hence  $N(u)$  is a triangle, and together with  $u$  we obtain a  $K_4$  as a subgraph of  $G$ , a contradiction.

For  $r = 5$ , by Theorem 3.1  $e \leq 3n - 6$  hence there is a vertex  $u \in G$  with degree  $d(u) \leq 5$ . Also  $d(u) \geq 4$  (as we may assume that  $u$  is not an isolated vertex). If  $d(u) = 4$  then the induced subgraph on  $\{u\} \cup N(u)$  is

$K_5$ , a contradiction. Otherwise,  $d(u) = 5$ . Every  $v \in N(u)$  has degree at least 3 in  $N(u)$ , hence  $e(N(u)) \geq \lceil 3 \cdot 5/2 \rceil = 8$ . But  $K_4 \not\prec N(u)$ , hence  $e(N(u)) \leq 2 \cdot 5 - 3 = 7$ , a contradiction. ■

**Proposition 3.3** *If  $G$  has an edge and each edge belongs to at least 4 triangles, then either  $G$  has a  $K_6$  minor, or  $G$  is a clique sum over  $K_r$  for some  $r \leq 4$  (i.e.  $G = G_1 \cup G_2, G_1 \cap G_2 = K_r, G_i \neq K_r, i = 1, 2$ ).*

*Proof:* We proceed as in the proof of Proposition 3.2: Assume that  $K_6 \not\prec G$ . W.l.o.g.  $G$  is connected. By Theorem 3.1  $e \leq 4n - 10$  hence there is a vertex  $u \in G$  with degree  $d(u) \leq 7$ , also  $d(u) \geq 5$ . If  $d(u) = 5$  then  $N(u) = K_5$ , a contradiction. Actually,  $N(u)$  is planar: since  $N(u)$  has at most 7 vertices, each of degree at least 4, if  $N(u)$  were not 4-connected, it must have exactly 7 vertices and two disjoint edges such that each of their 4 vertices is adjacent to the remaining 3 vertices of  $N(u)$  (whose removal disconnect  $N(u)$ ); but such graph has a  $K_5$  minor. As  $K_5 \not\prec N(u)$ ,  $N(u)$  is 4-connected. Now Wagner's structure theorem for  $K_5$ -minor free graphs ([5], Theorem 8.3.4) asserts that  $N(u)$  is planar.

If  $d(u) = 6$ , then  $12 = 3 \cdot 6 - 6 \geq e(N(u)) \geq 4 \cdot 6/2 = 12$  hence  $N(u)$  is a triangulation of the 2-sphere  $S^2$ . If  $d(u) = 7$ , then  $15 = 3 \cdot 7 - 6 \geq e(N(u)) \geq 4 \cdot 7/2 = 14$ . We will show now that  $N(u)$  cannot have 14 edges, hence it is a triangulation of  $S^2$ : Assume that  $N(u)$  has 14 edges, so each of its vertices has degree 4, and  $N(u)$  is a triangulation of  $S^2$  minus an edge. Let us look to the unique square (in a planar embedding) and denote its vertices by  $A$ . The number of edges between  $A$  and  $N(u) - A$  is 8. Together with the 4 edges in the subgraph induced by  $A$ , leaves two edges for the subgraph induced by  $N(u) - A = \{a, b, c\}$ ; let  $a$  be their common vertex. We now look at the neighborhood of  $a$  in a planar embedding (it is a 4-cycle):  $b, c$  must be opposite in this square as  $\{b, c\}$  is missing. Hence for  $v \in A \cap N(a)$  we get that  $v$  has degree 5, a contradiction.

Now we are left to deal with the case where  $N(u)$  is a triangulation of  $S^2$ , and hence a maximal  $K_5$ -minor free graph. If  $G$  is the cone over  $N(u)$  with apex  $u$ , then every edge in  $N(u)$  belongs to at least 3 triangles in  $N(u)$ . By Proposition 3.2,  $N(u)$  has a  $K_5$  minor, a contradiction. Hence there exists a vertex  $w \neq u, w \in G \setminus N(u)$ . Denote by  $[w]$  the set of all vertices in  $G$  connected to  $w$  by a path disjoint from  $N(u)$ . Denote by  $N'(w)$  the induced graph on the vertices in  $N(u)$  that are neighbors of some vertex in  $[w]$ . If  $N'(w)$  is not a clique, there are two non-neighbors  $x, y \in N'(w)$ , and a path through vertices of  $[w]$  connecting them. This path together with the cone over  $N(u)$  with apex  $u$  form a subgraph of  $G$  with a  $K_6$  minor, a contradiction.

Suppose  $N'(w)$  is a clique (it has at most 4 vertices, as  $N(u)$  is planar). Then  $G$  is a clique sum of two graphs that strictly contain  $N'(w)$ : Let  $G_1$  be the induced graph on  $[w] \cup N'(w)$  and let  $G_2$  be the induced graph on

$G \setminus [w]$ . Then  $G = G_1 \cup G_2$  and  $G_1 \cap G_2 = N'(w)$ . ■

**Remark:** In view of Theorem 3.1 for the case  $r = 7$ , we may expect the following to be true:

**Problem 3.4** *If  $G$  has an edge and each edge belongs to at least 5 triangles, then either  $G$  has a  $K_7$  minor, or  $G$  is a clique sum over  $K_l$  for some  $l \leq 6$ .*

If true, it extends the assertion of Theorem 1.2 to the case  $r = 7$ . By now we can show only the weaker assertion

$$G \text{ has a generic } 5\text{-stress} \Rightarrow K_7^- \prec G,$$

by using similar arguments to those used for proving Theorem 1.2 ( $K_7^-$  is  $K_7$  minus an edge).

## 4 Proof of Theorem 1.2

For  $r = 2$  the assertion of the theorem is trivial. Suppose  $K_r \not\prec G$ , and contract edges belonging to at most  $r - 3$  triangles as long as it is possible. Denote the resulted graph by  $G'$ . Repeated application of Theorem 2.2 asserts that if  $G'$  is generically  $(r - 2)$ -stress free, then so is  $G$ . In case  $G'$  has no edges, it is trivially  $(r - 2)$ -stress free. Otherwise,  $G'$  has an edge, and each edge belongs to at least  $r - 2$  triangles. For  $2 < r < 6$ , by Proposition 3.2  $G'$  has a  $K_r$  minor, hence so has  $G$ , a contradiction. For  $r = 6$ , by Proposition 3.3  $G'$  either has a  $K_6$  minor which leads to a contradiction, or  $G'$  is a clique sum over  $K_r$  for some  $r \leq 4$ . In the later case, denote  $G' = G_1 \cup G_2$ ,  $G_1 \cap G_2 = K_r$ . As the graph of a simplex is  $k$ -rigid for any  $k$ , by Theorem 2.1 it is enough to show that each  $G_i$  is generically  $(r - 2)$ -stress free, which follows from induction hypothesis on the number of vertices. ■

**Remark:** We can prove the case  $r = 5$  avoiding Mader's theorem, by using Wagner's structure theorem for  $K_5$ -minor free graphs ([5], Theorem 8.3.4) and Theorem 2.1. Using Wagner's structure theorem for  $K_{3,3}$ -minor free graphs ([5], ex.20 on p.185) and Theorem 2.1, we conclude that  $K_{3,3}$ -minor free graphs are generically 4-stress free.

## Acknowledgements

I would like to thank my advisor prof. Gil Kalai for many helpful discussions, and prof. Carsten Thomassen for his contribution to the Minors Section.

## References

- [1] L. Asimov and B. Roth, The rigidity of graphs, *Trans. Amer. Math. Soc.*, **245** (1978), 279-289.

- [2] L. Asimov and B. Roth, The rigidity of graphs: part II, *J. Math. Anal. Appl.*, **68** (1979), 171-190.
- [3] L. Cauchy, Sur les polygones et les polyèdres, *Second Memoire, I. Ecole Polytechnique*, **245** (1813) (=Oeuvres complètes d'Augustin Cauchy, 2nd sér., Tome 1 (1905), pp. 26-38).
- [4] Y. Colin de Verdière, Sur un nouvel invariant des graphes et un critère de planarité, *J. Combi. Th. Ser. B*, **50** (1990), 11-21. (English translation: On a new graph invariant and a criterion for planarity, in Graph Structure Theory (N. Robertson and P. Seymour eds.), *Contemp. Math.*, A.M.S., (1993), pp. 137-147.)
- [5] R. Diestel, *Graph Theory*, Second edition, Springer-Verlag, N.Y., (2000).
- [6] H. Gluck, Almost all simply connected closed surfaces are rigid, in Geometric topology. Lecture Notes in Math., Vol.438, pp.225-239 (1975).
- [7] G. Kalai, Rigidity and the lower bound theorem, *Inven. Math.*, **88**, (1987), 125-151.
- [8] G. Kalai, Algebraic Shifting, *Advanced Studies in Pure Math.*, **33** (2002), 121-163.
- [9] K. Kuratowski, Sur le problème des courbes gauches en topologie, *Fund. Math.*, **15**, (1930), 271-283.
- [10] K. W. Lee, Generalized stress and motion, in Polytopes: Abstract, Convex and Computational (T. Briztriczky et al., eds.), (1995) 249-271.
- [11] L. Lovász and A. Schrijver, A Borsuk theorem for antipodal links and a spectral characterization of linklessly embeddable graphs, *Proc. Amer. Math. Soc.*, **126** (1998), 1275-1285.
- [12] W. Mader, Homomorphiesätze für Graphen, *Math. Ann.* **178**, (1968), 154-168.
- [13] W. Mader,  $3n - 5$  edges do force a subdivision of  $K_5$ , *Combinatorica* Vol.**18**, **4**, (1998), 569-595.
- [14] E. Nevo, Embeddability and stresses of graphs, arXiv: math.CO/0411009.
- [15] N. Robertson, P.D. Seymour and R. Thomas, Linkless embeddings of graphs in 3-space, *Bull. Amer. Math. Soc.*, **28**, (1993), 84-89.

- [16] Z. Song, The extremal function for  $K_8^-$  minors, submitted, [www.math.gatech.edu/~song/K8.pdf](http://www.math.gatech.edu/~song/K8.pdf).
- [17] E. Steinitz and H. Rademacher, Volgesungen über die Theorie der Polyeder. Berlin-Göttingen Springer 1934.
- [18] W. Whiteley, Vertex splitting in isostatic frameworks, *Struc. Top.*, **16** (1989), 23-30.